

Generalized theta functions for holomorphic triples

Dissertation

zur Erlangung des akademischen Grades eines
Doktors der Naturwissenschaften (Dr.rer.nat.)

von

Quyet Thang Truong

geboren in Vietnam

vorgelegt beim Fachbereich Mathematik
der Universität Duisburg-Essen
Campus Essen

Essen 2015

Betreuer

Prof. Dr. Georg Hein

Gutachter

Prof. Dr. Georg Hein

Prof. Dr. Alexander Schmitt

Datum der mündlichen Prüfung

07. Juli 2015

Vorsitz

Prof. Dr. Jochen Heinloth

Contents

Introduction	iii
Acknowledgements	vii
1 Vector bundles on curves	1
1.1 Semistability of vector bundles	1
1.2 The Harder-Narasimhan filtration for vector bundles	3
2 Bogomolov's inequality for vector bundles on $X \times \mathbb{P}^1$ in case of arbitrary characteristic	9
2.1 Numerical equivalence of divisors on ruled surfaces	9
2.2 Bogomolov's inequality	10
2.3 Restriction theorems	16
3 Triples and their moduli spaces	23
3.1 Triples and their stability	23
3.2 Constraints on the parameter α	26
3.3 The corresponding extensions on $X \times \mathbb{P}^1$ of triples	28
3.3.1 The construction of extensions	28
3.3.2 $\mathrm{SL}(2)$ -invariant subbundles	29
3.4 Moduli space of triples	33
4 Generalized theta line bundles	35
4.1 Theta divisors for vector bundles on curves	35
4.1.1 Determinant line bundles	35
4.1.2 Generalized theta functions and generalized theta divisors	36
4.1.3 Pluritheta linear series on $U_X(r, d)$	39

4.2	Determinant line bundles and invariant sections for triples	40
4.2.1	Triple product	40
4.2.2	α -orthogonal triples	40
4.2.3	Determinant line bundles for triples	44
4.2.4	Invariant sections	46
4.3	Base point freeness	47
4.3.1	Generalized theta line bundles for triples	47
4.3.2	Linear systems	49
4.3.3	Corollaries	50
4.4	Langton's valuative criterion for triples	54
4.4.1	Elementary transformations	54
4.4.2	Valuative criterion	56

Bibliography	59
---------------------	-----------

Introduction

Let X be a smooth projective curve over an algebraically closed field k of arbitrary characteristic. A triple on X is a triple (E_2, E_1, φ) where E_1, E_2 are vector bundles on X and $\varphi : E_2 \rightarrow E_1$ is a morphism. For triples, we can introduce the notion of (semi)stability, depending on a real parameter α , and then study the moduli space of α -(semi)stable triples. The properties of triples and their moduli spaces have been investigated in [2] by Bradlow and García-Prada, in [3] by the same authors with Gothen, in [17] by Schmitt in characteristic 0 or in [1] by Álvarez-Cónsul in case of arbitrary characteristic, etc. Particularly, in [2], Bradlow and García-Prada showed that there is a one-to-one correspondence between triples $T = (E_2, E_1, \varphi)$ and the extensions $E_T \in \text{Ext}^1(p^*E_2, p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2))$ on $X \times \mathbb{P}^1$ defined by T . By showing that T is α -(semi)stable if and only if E_T is $H(\alpha)$ -(semi)stable where $H(\alpha)$ is an ample divisor on $X \times \mathbb{P}^1$, Bradlow and García-Prada constructed the moduli space of α -(semi)stable triples as a closed subscheme of the moduli space of semistable vector bundles of fixed type on $X \times \mathbb{P}^1$. Let $C \in |mH(\alpha)|$ be a smooth projective curve on $X \times \mathbb{P}^1$. It is well known that for m large enough, the restriction of E_T to C is still semistable. We then obtain a morphism from moduli space of α -semistable triples to the moduli space of semistable vector bundles on C which allows us to study triples and their moduli space from moduli space of vector bundles on C .

Let $U_C(r, d)$ be the moduli space of semistable vector bundles of rank r and degree d on C . For a given vector bundle F on C such that $\chi(E \otimes F) = 0$ for some $E \in U_C(r, d)$ (hence for all $E \in U_C(r, d)$), we set $\Theta_F = \{E \in U_C(r, d) | H^0(C, E \otimes F) \neq 0\}$. It was proven by Drezet and Narasimhan in [5] that Θ_F is the underlying space of an effective Cartier divisor, called a generalized theta divisor, if it is a proper subset of $U_C(r, d)$. Moreover, for any vector bundle F' such that $[F'] = n[F]$ in the Grothendieck's group of coherent sheaves $K(C)$, we have $\Theta_{F'} \in |n\Theta_F|$. On smooth projective curve C , it was first proven by Faltings in [8] and then by Seshadri in [18] that a vector bundle E is semistable if and only if there exists a vector bundle F such that $E \otimes F$ is cohomologically trivial, i.e. $H^*(X, E \otimes F) = 0$. We say that F is orthogonal to E . It follows that Θ_F is a generalized theta divisor if F is orthogonal to some $E \in U_C(r, d)$. In this case, the line bundle $\theta_F = \mathcal{O}_{U_C(r, d)}(\Theta_F)$ has a section ζ_F , called a generalized theta function, such that its zero divisor is Θ_F . Studying properties of linear

system $|\Theta_F|$: base point freeness, separation properties, ampleness, global generatedness, etc, allows us to construct and study the moduli space of vector bundles on curves by a GIT-free way (see [8], [18], [11], [16], [7], [6], etc). As mentioned above, there is a morphism from moduli space of α -semistable triples on X to the moduli space of semistable vector bundles on C . So there should be a natural way to define and study theta line bundles and theta functions on the moduli space of α -semistable triples.

The aims of this thesis are to extend some properties of α -semistable triples on smooth projective curves to arbitrary characteristic and to add the missing pieces of the study of moduli space of triples: generalized theta line bundles and generalized theta functions. Studying properties of these objects also allows us to define the orthogonal triples and furthermore prepare the ingredients for a GIT-free construction of moduli space of triples.

More in details, we firstly recall in **Chapter 1** the definitions, basic properties of slope semistability and global generatedness of vector bundles on curves. In particular, we recall the existence of the Harder-Narasimhan filtration for vector bundles and also a modified Harder-Narasimhan filtration which will be used later.

In **Chapter 2**, we recall the numerical equivalence of divisors on ruled surfaces, typically on $X \times \mathbb{P}^1$. We then prove the following Bogomolov's inequality for vector bundles on $X \times \mathbb{P}^1$ in case of arbitrary characteristic,

Theorem 1 (Thm. 2.2.4). *Let E be a vector bundle on $X \times \mathbb{P}^1$ and H be an ample divisor. If E is H -semistable then*

$$\Delta(E) = (\text{rank}(E) - 1)c_1^2(E) - 2\text{rank}(E)c_2(E) \leq 0.$$

As a consequence of Bogomolov's inequality, we obtain then a restriction theorem for semistable vector bundles on $X \times \mathbb{P}^1$ (Cor. 2.3.4).

In **Chapter 3** we collect the definitions and basic properties of triples, α -semistability of triples and also the existence of the moduli space of triples. For each triple T , we construct a vector bundle E_T on $X \times \mathbb{P}^1$ (see Section 3.3). Let p, q be the projections from $X \times \mathbb{P}^1$ onto X and \mathbb{P}^1 , respectively. Let $\alpha = \frac{a}{b}$ be a positive rational number. Let us fix $H(\alpha) = aF_p + bF_q$ an ample divisor on $X \times \mathbb{P}^1$ where F_p, F_q are any p -fiber and q -fiber, respectively. We give then a proof of the following result of Bradlow and García-Prada in [2] which works in arbitrary characteristic.

Theorem 2 (Thm. 3.3.4). *The triple $T = (E_2, E_1, \varphi)$ is α -(semi)stable if and only if the extension E_T defined by T is $H(\alpha)$ -(semi)stable.*

Chapter 4 is the main part of the thesis where we define and study properties of generalized theta line bundles and generalized theta functions on moduli space of triples. Let $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ be a family of α -semistable triples on X parametrized by a scheme S . Using the construction of the determinant line bundle associated to a flat family of coherent

sheaves (see Section 4.1.1) and the triple product (see Section 4.2.1), we can define the determinant line bundle for triples (Def 4.2.5). For any triple $F = (F_2 \xrightarrow{\psi} F_1)$ on X , we then obtain a line bundle $\theta_{F_2 \rightarrow F_1}$ on S . In particular, if $\psi : F_2 \rightarrow F_1$ is surjective and the Euler characteristic of triple product $\mathcal{T}_s \times F$ is zero for all $s \in S$ then $\theta_{F_2 \rightarrow F_1}$ has a section $\zeta_{F_2 \rightarrow F_1}$ (see Section 4.2.4). Let $C \in |mH(\alpha)|$ be a smooth projective curve. As an application of Bogomolov's inequality, $E_T|_C$ is still semistable for any $m \geq m_0 = \lceil \frac{1-r}{r} \Delta(E_T) + 1 \rceil$ (Cor. 2.3.4). As $E_T|_C$ is semistable, the existence of orthogonal bundles F_C allows us to construct a α -orthogonal triple $F = (F_2 \xrightarrow{\psi} F_1)$ (Def 4.2.2) which characterizes the α -semistability of T :

Theorem 3 (Thm. 4.2.3). *Let $T = (E_2 \xrightarrow{\varphi} E_1)$ be a triple on X and $\alpha = \frac{a}{b} \in \mathbb{Q}$. Then T is α -semistable if and only if it has a α -orthogonal triple.*

Let us fix $C \in |m_0 H(\alpha)|$. Let \mathcal{M} be the moduli space of α -semistable triples of type (r_1, r_2, d_1, d_2) on X and $U_C(r, d)$ be the moduli space of semistable vector bundles on C of rank $r = r_1 + r_2$ and degree $d = m_0 \deg_{H(\alpha)}(E_T)$. We have a morphism

$$\mathfrak{t} : \mathcal{M} \longrightarrow U_C(r, d)$$

sending T to $E_T|_C$ (Lemma 4.3.1). We define the generalized theta line bundle and generalized theta functions to be the pullback by \mathfrak{t} of the corresponding ones on $U_C(r, d)$ (Def. 4.3.2). Let θ_{F_0} be a basic theta divisor on $U_C(r, d)$ and $|p\theta_{F_0}|$ be a pluritheta linear series. Let $\mathcal{L} = \mathfrak{t}^* \theta_{F_0}$ be a basic theta line bundle on \mathcal{M} . Then for each p , we have a generalized theta line bundle \mathcal{L}^p and a linear system $V(\mathcal{L}^p) \subset H^0(\mathcal{M}, \mathcal{L}^p)$ spanned by generalized theta functions. The first and most important applications which we have from the existence of α -orthogonal triples are the base point freeness, ampleness and the universal property of the generalized theta line bundle for triples. Let F_C be a vector bundle on C such that $[F_C] = p[F_0] \in K(C)$. Using F_C we construct a triple $F = (F_2 \xrightarrow{\psi} F_1)$ (see proof of Lemma 4.2.4 or (4.27)), such that the line bundle $\theta_{F_2 \rightarrow F_1}$ has a geometric section $\zeta_{F_2 \rightarrow F_1}$ whose zero divisor is defined by

$$\Theta_{F_2 \rightarrow F_1} = \{s \in S \mid F \text{ is not a } \alpha\text{-orthogonal triple of } \mathcal{T}_s\}$$

(see (4.21) and (4.25)). Let $\text{Comp}(p, F)$ be the set of all triples $F' = (F'_2 \xrightarrow{\psi'} F'_1)$ such that $[F'_i] = [F_i] \in K(X)$. Then all the line bundles $\theta_{F'_2 \rightarrow F'_1}$ are isomorphic to $\theta_{F_2 \rightarrow F_1}$ (see Lemma 4.2.6). Therefore we obtain on S a linear system $V(p, F) \subset H^0(S, \theta_{F_2 \rightarrow F_1})$ spanned by sections $\zeta_{F'_2 \rightarrow F'_1}$. By the result of Esteves and Popa on the base point freeness of pluritheta linear series in [7], we obtain the following base point freeness of linear systems $V(\mathcal{L}^p)$ and $V(p, F)$.

Corollary 4 (Cor. 4.3.3). *For any $p \geq r^2 + r$, then*

- (i) *The linear systems $V(\mathcal{L}^p)$ and $V(p, F)$ are base point free.*

(ii) *There exists a triple $F' = (F'_2 \xrightarrow{\psi'} F'_1) \in \text{Comp}(p, F)$ such that*

$$\overline{\Theta}_{F'_2 \rightarrow F'_1} = \{T \in \mathcal{M} \mid F' \text{ is not a } \alpha\text{-orthogonal triple of } T\}$$

is a Cartier divisor on \mathcal{M} and

$$\mathcal{L}^p \cong \mathcal{O}_{\mathcal{M}}(\overline{\Theta}_{F'_2 \rightarrow F'_1}).$$

Similar to the construction of extensions on $X \times \mathbb{P}^1$ defined by triples on X , for each family \mathcal{T} of α -semistable triples on X parametrized by S , we obtain an induced family $E_{\mathcal{T}}$ of $H(\alpha)$ -semistable vector bundles on $X \times \mathbb{P}^1$ and then a family $\mathcal{E} = E_{\mathcal{T}}|_C$ of semistable vector bundles on C parametrized by S . We then have a commutative diagram of induced morphisms

$$\begin{array}{ccc} S & \xrightarrow{f_{\mathcal{E}}} & U_C(r, d). \\ & \searrow f_{\mathcal{T}} & \nearrow \iota \\ & \mathcal{M} & \end{array}$$

By the universal property and the ampleness of the generalized theta line bundle θ_{F_C} on $U_C(r, d)$ (Thm. 4.1.2), we obtain the similar properties for \mathcal{L}^p .

Corollary 5 (Cor. 4.3.4). *Let $p \geq 1$ and $F'' = (F''_2 \xrightarrow{\psi''} F''_1)$ be any triple in $\text{Comp}(p, F)$. Then for any family $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ of α -semistable triples on X parametrized by S , the line bundle $\theta_{F''_2 \rightarrow F''_1}$ descends to \mathcal{L}^p by the induced morphism $f_{\mathcal{T}} : S \rightarrow \mathcal{M}$.*

Corollary 6 (Cor. 4.3.5). *The line bundle \mathcal{L} on \mathcal{M} is ample.*

Another important consequence of Theorem 3 is Langton's valuative criterion for triples. Let R be a discrete valuation ring and $X_{\eta} = \eta \times X$ be the generic fiber of the first projection

$$\text{Spec}(R) \times X \rightarrow \text{Spec}(R) = \{0, \eta\}.$$

Theorem 7 (Thm. 4.4.4). *Let T_{η} be a α -semistable triple of type (r_1, r_2, d_1, d_2) on the generic fiber X_{η} . Then there exists a triple \mathcal{T} on X_R such that $\mathcal{T}|_{X_{\eta}} \cong T_{\eta}$ and $T_0 := \mathcal{T}|_{X_0}$ is α -semistable.*

Acknowledgements

This work would not have been possible without the dedicated and scientific guide, ceaseless support and encouragement of my supervisor Prof. Dr. Georg Hein. I would like to express my warmest gratitude to him for all the best things he has instructed and done for me since I was a Master student.

I give my sincere thanks to the German Academic Exchange Service (DAAD) for supporting me during most of my PhD period. I have also received a lot of help from many people in “Essener Seminar für Algebraische und Arithmetik” at the Mathematics Department, University of Duisburg- Essen, to whom I am also grateful. Especially, I really need to give my thanks to Prof. Dr. Vytautas Paškūnas, who has always helped me in carrying out the procedure of my scholarship extension with DAAD.

Last but not least, I feel warmly thankful to my family and friends, who always stand by and encourage me.

Chapter 1

Vector bundles on curves

1.1 Semistability of vector bundles

Let X be a smooth projective curve over an algebraically closed field k of arbitrary characteristic. By a vector bundle on X , we mean a locally free coherent sheaf on X . Let E be a nonzero vector bundle on X , $d = \deg(E)$ and $r = \text{rank}(E)$. Then we define the slope of E to be $\mu(E) = \frac{d}{r}$.

Definition 1.1.1. *Let E be a nonzero vector bundle on X . Then E is called (semi)stable if for any nonzero subsheaf $E' \subsetneq E$ we have $\mu(E')(\leq)\mu(E)$.*

Here we use the notation (\leq) introduced by Huybrechts and Lehn in [12, p.11]: If the word “(semi)stable” and the relation sign “ (\leq) ” appear together in a statement, then this statement stands for two assertions, one with “stable” and strict inequality “ $<$ ” and the other with “semistable” and the mild inequality “ \leq ”.

Remark 1.1.2. Assume that

$$0 \longrightarrow F \longrightarrow E \longrightarrow G \longrightarrow 0$$

is an exact sequence of vector bundles. Then we have

$$\begin{aligned}\deg(E) &= \deg(F) + \deg(G), \\ \text{rank}(E) &= \text{rank}(F) + \text{rank}(G).\end{aligned}$$

It implies that

$$\mu(F)(\leq)\mu(E) \iff \mu(E)(\leq)\mu(G).$$

Hence we can also formulate semistability in terms of quotient bundles: E is semistable if and only if for any nonzero quotient G of E , we have $\mu(E) \leq \mu(G)$.

The following properties of semistable vector bundles can be seen easily by the definition.

Lemma 1.1.3. *Let E be a vector bundle on X and L be a line bundle. Then the following holds:*

(i) L is stable.

(ii) E is (semi)stable if and only if $E \otimes L$ is (semi)stable.

(iii) E is (semi)stable if and only if E^\vee , its dual bundle, is (semi)stable.

Proposition 1.1.4. *Let E and F be vector bundles on X .*

(i) *If E and F are both semistable and $\mu(E) > \mu(F)$ then $\text{Hom}_X(E, F) = 0$.*

(ii) *If E is stable then it is simple, i.e. $\text{End}(E) \cong k$.*

Proof. Let $\phi \in \text{Hom}(E, F)$ be a morphism. If $\phi \neq 0$ then $I = \text{Im } \phi \subset F$ is not only a nonzero subsheaf of F but also a quotient of E . Since E and F are semistable, we have

$$\mu(E) \leq \mu(I) \leq \mu(F).$$

This proves (i). For (ii), we show that any endomorphism of E is in form λId_E for some $\lambda \in k$. Let $\phi \in \text{End}(E)$ be a nontrivial morphism and I be the image of ϕ . As we have shown above, $\mu(E) < \mu(I) < \mu(E)$ if $I \subsetneq E$. This is impossible. Hence ϕ must be surjective and hence an isomorphism. If $\phi \neq \lambda \text{Id}_E$ then there exists $x \in X(k)$ such that $\phi_x \in \text{End}_k(E \otimes k(x))$ is not in form λId . Let λ be a eigentvalue of ϕ_x . Then $\phi - \lambda \text{Id}_E \in \text{End}(E)$ is nontrivial but not surjective. Hence it is not an isomorphism. This gives us a contradiction. \square

For any coherent sheaf E on X , $\chi(E)$ is the Euler characteristic of E , defined by

$$\chi(E) = \dim H^0(X, E) - \dim H^1(X, E).$$

Theorem 1.1.5 (Riemann-Roch formula - [9, Thm. 4.1, p.432]). *Let E be a coherent sheaf on smooth projective curve X of genus g_X . Then*

$$\chi(E) = \deg(E) + \text{rank}(E)(1 - g_X).$$

It follows from Riemann-Roch formula that

$$\mu(E) = \frac{\chi(E)}{\text{rank}(E)} + g_X - 1$$

for any nontrivial vector bundle E on X .

Proposition 1.1.6. *Let E and F be vector bundles on X . If $F \neq 0$ and $H^*(E \otimes F) = 0$, i.e. $H^0(X, E) = H^1(X, E) = 0$, then E is semistable.*

Proof. Assume that $F \neq 0$ and $H^*(X, E \otimes F) = 0$. Since $H^*(X, E \otimes F) = 0$, $\chi(E \otimes F) = 0$. It implies that $g_X - 1 = \mu(E \otimes F)$. If E is not semistable then there exists a nontrivial subbundle $G \subsetneq E$ such that $\mu(G) > \mu(E)$. It follows that

$$\mu(G \otimes F) > \mu(E \otimes F) = g_X - 1$$

and hence $\chi(G \otimes F) > 0$. But we also have

$$\chi(G \otimes F) = \dim H^0(X, G \otimes F) - \dim H^1(X, G \otimes F) \leq 0$$

since $H^0(X, G \otimes F) \subseteq H^0(X, E \otimes F) = 0$. This is a contradiction. So E is semistable. \square

Definition 1.1.7. *A vector bundle E on X is called globally generated if the evaluation map*

$$e : H^0(X, E) \otimes \mathcal{O}_X \longrightarrow E$$

is surjective.

Theorem 1.1.8 ([9, Thm. 5.17, p.121; Thm. 5.2, p.228]). *Let E be a vector bundle on X . Then there exists a number $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$,*

(i) *$E(n)$ is globally generated.*

(ii) *$H^q(X, E(n)) = 0$ for all $q > 0$.*

The following properties of globally generated vector bundles are useful later.

Proposition 1.1.9 ([11, Prop. 2.6]). *Let E be a globally generated vector bundle of rank r on X . Then we have the following exact sequences*

$$0 \longrightarrow \mathcal{O}_X^{r-1} \longrightarrow E \longrightarrow \det(E) \longrightarrow 0$$

and

$$0 \longrightarrow \det(E)^{-1} \longrightarrow \mathcal{O}_X^{r+1} \longrightarrow E \longrightarrow 0.$$

1.2 The Harder-Narasimhan filtration for vector bundles

In this section we will explain the existence and some elementary properties of the Harder-Narasimhan filtration for vector bundles on a smooth projective curve. Moreover, we can make a slight modification on the index set of the Harder-Narasimhan filtration so that it can be indexed by all of rational numbers.

Definition 1.2.1. *Let E be a vector bundle over a smooth projective curve X . A Harder-Narasimhan filtration for E is a flag*

$$0 = E_0 \subset E_1 \subset E_2 \subset \dots \subset E_k = E$$

of subbundles of E having the following two properties:

- (i) E_i/E_{i-1} is semistable for $i = 1, \dots, k$,
- (ii) $\mu(E_{i+1}/E_i) < \mu(E_i/E_{i-1})$ for $i = 1, \dots, k-1$.

By Riemann-Roch formula,

$$\mu(E') \leq g_X - 1 + \dim H^0(X, E')$$

for any nonzero subsheaf E' of E . Hence there exists a subsheaf of E which is maximal among the subsheaves of E of maximal slope, i.e. there exists a subsheaf $F \subseteq E$ such that for all $G \subseteq E$, $\mu(G) \leq \mu(F)$ and in case of equality, $G \subseteq F$. In fact, F is semistable and unique. We call it the maximal destabilizing subsheaf. The existence of the maximal destabilizing subsheaf induces the existence of the Harder-Narasimhan filtration for a vector bundle on X .

Theorem 1.2.2 ([12, Thm. 1.3.4]). *Let E be a vector bundle over a smooth projective curve X , then there exists a unique Harder-Narasimhan filtration for E .*

The following properties come directly from the definition of the Harder-Narasimhan filtration.

Proposition 1.2.3. *Let E be a vector bundle with the Harder-Narasimhan filtration*

$$0 = E_0 \subset E_1 \subset E_2 \subset \dots \subset E_k = E.$$

Then we have

- (i) $\mu(E_j/E_i) > \mu(E_t/E_i)$ for $i < j < t$.
- (ii) $\mu(E_t/E_i) > \mu(E_t/E_j)$ for $i < j < t$.
- (iii) $\mu(E_1) > \mu(E_2) > \dots > \mu(E_k) = \mu(E)$.

Let us set $\mu_i = \mu(E_i/E_{i-1})$ for $i = 2, \dots, k-1$ and $\mu_{\max} = \mu(E_1)$, $\mu_{\min} = \mu(E/E_{k-1})$. By a slight modification, we can index the Harder-Narasimhan filtration by all of rational numbers as follows: For each $\rho \in \mathbb{Q}$, we set $E_{(\rho)} = E_i$ where i is the index such that

$$\mu_i \geq \rho > \mu_{i+1}.$$

It follows that

$$E_{(\rho)} = 0 \iff \rho > \mu_{\max} \text{ and } E_{(\rho)} = E \iff \rho \leq \mu_{\min}. \quad (1.1)$$

Proposition 1.2.4. *The following properties are true:*

(i) *If $\rho > \tau$ then $E_{(\rho)} \subseteq E_{(\tau)}$. If $E_{(\rho)} \subsetneq E_{(\tau)}$ then $\rho > \tau$.*

(ii) *$\mu_{\min}(E_{(\tau)}) \geq \tau, \mu_{\max}(E/E_{(\tau)}) < \tau$.*

(iii) *$(E_{(\rho)})_{(\tau)} = (E_{(\tau)})_{(\rho)} = E_{(\max\{\rho, \tau\})}$.*

Proof. Assume that E has the following Harder-Narasimhan filtration

$$0 = E_0 \subset E_1 \subset E_2 \subset \dots \subset E_k = E.$$

Assume $\rho > \tau$ and $E_{(\tau)} = E_i$. Then $\mu_i \geq \tau > \mu_{i+1}$. It implies that

$$\rho > \mu_i \text{ or } \mu_i \geq \rho > \mu_{i+1}.$$

By definition, $E_{(\rho)} \subseteq E_i = E_{(\tau)}$. If $E_{(\rho)} \subsetneq E_i$ then $E_{(\rho)} \subseteq E_{i-1}$. Therefore

$$\rho \geq \mu_{i-1} > \tau$$

and we have (i).

As $E_{(\tau)} = E_i$, the Harder-Narasimhan filtration of $E_{(\tau)}$ is induced from the one of E ,

$$0 = E_0 \subset E_1 \subset E_2 \subset \dots \subset E_i.$$

By definition, we have

$$\mu_{i+1} < \tau \leq \mu_i = \mu(E_i/E_{i-1}) = \mu_{\min}(E_{(\tau)}).$$

Similarly,

$$0 = E_i/E_i \subset E_{i+1}/E_i \subset E_{i+2}/E_i \subset \dots \subset E_k/E_i = E/E_i$$

is the Harder-Narasimhan filtration of $E/E_{(\tau)}$. Then

$$\mu_{\max}(E/E_{(\tau)}) = \mu(E_{i+1}/E_i) = \mu_{i+1} < \tau$$

and we get (ii).

If $\rho \leq \tau$ then $(E_{(\tau)})_{(\rho)} = E_{(\tau)}$ since $\rho \leq \mu_{\min}(E_{(\tau)})$ by (1.1) and (ii). If $\rho > \tau$ then

$$E_{(\rho)} = E_{(\max\{\rho, \tau\})} \subseteq E_{(\tau)} \subseteq E$$

by (i). Therefore $E_{(\rho)} = (E_{(\rho)})_{(\rho)} \subseteq (E_{(\tau)})_{(\rho)} \subseteq E_{(\rho)}$. It proves (iii). \square

Corollary 1.2.5. *Let $f : E \longrightarrow F$ be a morphism of vector bundles on X . Then for every $\rho \in \mathbb{Q}$, there exists a unique morphism*

$$f_\rho : E_{(\rho)} \longrightarrow F_{(\rho)}$$

making the following diagram commutative

$$\begin{array}{ccc} E_{(\rho)} & \xrightarrow{f_\rho} & F_{(\rho)} \\ \downarrow & & \downarrow \\ E & \xrightarrow{f} & F \end{array}$$

Proof. Assume that E and F have the Harder-Narasimhan filtration

$$0 = E_0 \subset E_1 \subset E_2 \subset \dots \subset E_k = E$$

and

$$0 = F_0 \subset F_1 \subset F_2 \subset \dots \subset F_{k'} = F,$$

respectively. Let $\rho \in \mathbb{Q}$ and $E_{(\rho)} = E_i \subseteq E$. The claim is trivial if $f(E_{(\rho)}) = 0$. Assume that $f(E_{(\rho)}) \neq 0$. Let j be the minimal index such that $f(E_{(\rho)}) \subseteq F_j$. Let $\tau = \mu(F_j/F_{j-1})$, then $F_{(\tau)} = F_j$ and the composition

$$E_i \xrightarrow{f} F_j \xrightarrow{\pi} F_j/F_{j-1}$$

is not zero. Suppose that $F_{(\tau)} \not\subseteq F_{(\rho)}$, then $\rho > \tau$. By definition, F_j/F_{j-1} is semistable with slope τ and for $1 \leq \nu \leq i$, $E_\nu/E_{\nu-1}$ is semistable with slope $\mu_\nu \geq \rho > \tau$. It is implied by Proposition 1.1.4 that $\text{Hom}(E_\nu/E_{\nu-1}, F_j/F_{j-1}) = 0$ for all $1 \leq \nu \leq i$. It follows that $\text{Hom}(E_i, F_j/F_{j-1}) = 0$. This is a contradiction since $\pi \circ f$ is not zero. \square

For each $\tau \in \mathbb{Q}$, we define the quotient

$$\text{gr}_\tau(E) := E_{(\tau)} / (\cup_{\rho > \tau} E_{(\rho)}).$$

It is clear that $\text{gr}_\tau(E) \neq 0$ if and only if $\tau = \mu(E_i/E_{i-1})$ for some $i \neq 0$. In that case,

$$\text{gr}_{\mu(E_i/E_{i-1})}(E) = E_i/E_{i-1}.$$

The following lemma will be used later.

Lemma 1.2.6. *Let E and F be two vector bundles on X with $\mu(E) > \mu(F)$. Then there exists a rational number τ such that*

$$\frac{\text{rank}(E_{(\tau)})}{\text{rank}(E)} > \frac{\text{rank}(F_{(\tau)})}{\text{rank}(F)}.$$

Proof. We set $M = \{\tau \in \mathbb{Q} \mid \text{rank}(\text{gr}_\tau(E)) + \text{rank}(\text{gr}_\tau(F)) > 0\}$. Then M is a finite set and we could write $M = \{\tau_1, \tau_2, \dots, \tau_n\}$ with $\tau_i > \tau_{i+1}$. We define two functions $\text{rank}_E, \text{rank}_F$ from M to \mathbb{N} by setting $\text{rank}_E(\tau_i) = \text{rank}(\text{gr}_{\tau_i}(E))$ and $\text{rank}_F(\tau_i) = \text{rank}(\text{gr}_{\tau_i}(F))$, respectively. Now we have

$$\begin{aligned} \mu(E) &= \frac{\deg(E)}{\text{rank}(E)} = \frac{\sum_{i=1}^n \deg(\text{gr}_{\tau_i}(E))}{\text{rank}(E)} \\ &= \sum_{i=1}^n \tau_i \frac{\text{rank}_E(\tau_i)}{\text{rank}(E)} = \sum_{i=1}^n (\tau_i - \tau_{i+1}) \sum_{j=1}^i \frac{\text{rank}_E(\tau_j)}{\text{rank}(E)}, \end{aligned}$$

where we set $\tau_{n+1} := 0$. Therefore we obtain

$$\mu(E) = \sum_{i=1}^{n-1} (\tau_i - \tau_{i+1}) \sum_{j=1}^i \frac{\text{rank}_E(\tau_j)}{\text{rank}(E)} + \tau_n.$$

The same formula holds for the slope of F . Since $\mu(E) > \mu(F)$,

$$\sum_{i=1}^{n-1} (\tau_i - \tau_{i+1}) \left(\sum_{j=1}^i \frac{\text{rank}_E(\tau_j)}{\text{rank}(E)} - \sum_{j=1}^i \frac{\text{rank}_F(\tau_j)}{\text{rank}(F)} \right) > 0.$$

We have $\tau_i > \tau_{i+1}$ for all i . It implies that there exists i such that

$$\sum_{j=1}^i \frac{\text{rank}_E(\tau_j)}{\text{rank}(E)} - \sum_{j=1}^i \frac{\text{rank}_F(\tau_j)}{\text{rank}(F)} > 0 \Leftrightarrow \frac{\text{rank}(E_{(\tau_i)})}{\text{rank}(E)} > \frac{\text{rank}(F_{(\tau_i)})}{\text{rank}(F)}.$$

□

Chapter 2

Bogomolov's inequality for vector bundles on $X \times \mathbb{P}^1$ in case of arbitrary characteristic

2.1 Numerical equivalence of divisors on ruled surfaces

Definition 2.1.1. *A geometrically ruled surface is a surface Y , together with a surjective morphism $\pi : Y \rightarrow X$ to a smooth curve X such that the fiber Y_x is isomorphic to \mathbb{P}^1 for every point $x \in X$ and π admits a section.*

One of the simplest examples of ruled surfaces which is our main subject is the ruled surface $X \times \mathbb{P}^1$ together with the first projection $p : X \times \mathbb{P}^1 \rightarrow X$, where X is a smooth projective curve.

Definition 2.1.2. *A divisor D on a surface Y is numerically equivalent to zero, written $D \equiv 0$, if the intersection number $D.E$ is zero for any divisor E . We say that D and E are numerically equivalent, written $D \equiv E$, if $D - E \equiv 0$.*

Let $\text{Num } Y$ be the quotient of $\text{Pic } Y$ by the subgroup of divisor classes numerically equivalent to zero. Then $\text{Num } Y$ is a finitely generated abelian group. The intersection pairing induces a nondegenerate bilinear pairing $\text{Num } Y \times \text{Num } Y \rightarrow \mathbb{Z}$.

Proposition 2.1.3 ([9, Prop. 2.3, p. 370]). *Let $\pi : Y \rightarrow X$ be a ruled surface, X_0 be a section and f be any fiber. Then*

$$\text{Pic } Y \cong \mathbb{Z} \oplus \pi^* \text{Pic } X$$

where \mathbb{Z} is generated by X_0 . Also

$$\text{Num } Y \cong \mathbb{Z} \oplus \mathbb{Z},$$

generated by X_0, f and satisfying $X_0.f = 1, f^2 = 0$.

On our ruled surface $X \times \mathbb{P}^1$, let $p : X \times \mathbb{P}^1 \longrightarrow X$ and $q : X \times \mathbb{P}^1 \longrightarrow \mathbb{P}^1$ be the projections. We can choose X_0 to be any q -fiber F_q and f be any p -fiber F_p . It implies that for any divisor class $D \in \text{Num}(X \times \mathbb{P}^1)$, we have $D \equiv aF_p + bF_q$ for some $a, b \in \mathbb{Z}$ where $F_p^2 = F_q^2 = 0, F_p.F_q = 1$. In particular, we have

Proposition 2.1.4 ([9, Prop. 2.20, p. 382]). *A divisor $H \equiv aF_p + bF_q$ on $X \times \mathbb{P}^1$ is ample if and only if $a, b \in \mathbb{Z}_{>0}$.*

Let $H \equiv aF_p + bF_q$ be an ample divisor on $X \times \mathbb{P}^1$, $D \equiv cF_p + dF_q$. Then $H^2 = 2ab > 0$. If $D.H = 0$ then $ad + bc = 0$. It implies that

$$D^2 = 2cd \leq 0. \quad (2.1)$$

We have $D^2H^2 = 4abcd$, $(D.H)^2 = (ad + bc)^2$. Hence

$$(D.H)^2 - D^2H^2 = (ad - bc)^2 \geq 0 \quad (2.2)$$

for any divisor D .

2.2 Bogomolov's inequality

In this section, we will prove Bogomolov's inequality for torsion free coherent sheaves on the ruled surface $X \times \mathbb{P}^1$ in case of arbitrary characteristic. Firstly, we recall the Grothendieck's splitting theorem for vector bundles on \mathbb{P}^1 .

Theorem 2.2.1 (Grothendieck's splitting theorem - [12, Thm. 1.3.1]). *Let E be a vector bundle of rank r on \mathbb{P}^1 . Then there exists a unique r -tuple $(a_1, a_2, \dots, a_r) \in \mathbb{Z}^r$ with $a_1 \leq a_2 \leq \dots \leq a_r$, such that*

$$E \cong \bigoplus_{i=1}^r \mathcal{O}_{\mathbb{P}^1}(a_i).$$

Recall that $p : X \times \mathbb{P}^1 \longrightarrow X$ and $q : X \times \mathbb{P}^1 \longrightarrow \mathbb{P}^1$ are the projections.

Lemma 2.2.2. *Let E be a vector bundle of rank r on $X \times \mathbb{P}^1$. Suppose that for any p -fiber F_p we have $E|_{F_p} \cong \mathcal{O}_{\mathbb{P}^1}^{\oplus r}$. Then there exists a vector bundle F on X such that $E \cong p^*F$.*

Proof. See [18, Lm. 4.1]. □

Proposition 2.2.3. *Let E be a vector bundle on $X \times \mathbb{P}^1$. Suppose that*

$$R^1p_*(E \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)) = 0.$$

Then there exist vector bundles E_0 and E_1 on X such that the following sequence of vector bundles on $X \times \mathbb{P}^1$

$$0 \longrightarrow p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \longrightarrow p^*E_0 \longrightarrow E \longrightarrow 0 \quad (2.3)$$

is exact.

Proof. Let $x \in X$ be any closed point and E be a vector bundle on $X \times \mathbb{P}^1$ such that $R^1p_*(E \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)) = 0$. Then E_x , the restriction of E on the fiber of p over x , is a vector bundle on $F_p = x \times \mathbb{P}^1 \cong \mathbb{P}^1$. We have

$$R^1p_*(E \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)) \otimes k(x) \cong H^1(\mathbb{P}^1, E_x(-1)).$$

So the condition $R^1p_*(E \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)) = 0$ implies that

$$H^1(\mathbb{P}^1, E_x(-1)) = 0$$

for all $x \in X$. On $\mathbb{P}^1 \times \mathbb{P}^1$ we have the following resolution of diagonal

$$0 \longrightarrow p_1^*\mathcal{O}_{\mathbb{P}^1}(-1) \otimes p_2^*\mathcal{O}_{\mathbb{P}^1}(-1) \longrightarrow \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1} \longrightarrow \mathcal{O}_{\Delta_{\mathbb{P}^1}} \longrightarrow 0, \quad (2.4)$$

where p_1, p_2 are the first and second projection from $\mathbb{P}^1 \times \mathbb{P}^1$. Consider the following commutative diagram

$$\begin{array}{ccccc} & & X \times \mathbb{P}^1 & \xrightarrow{q} & \mathbb{P}^1 \\ & \nearrow p & \uparrow p_{13} & & \uparrow p_2 \\ X & & X \times \mathbb{P}^1 \times \mathbb{P}^1 & \xrightarrow{p_{23}} & \mathbb{P}^1 \times \mathbb{P}^1 \\ & \nwarrow p & \downarrow p_{12} & & \downarrow p_1 \\ & & X \times \mathbb{P}^1 & \xrightarrow{q} & \mathbb{P}^1 \end{array}$$

where p_{ij} are the projections. Pulling back the exact sequence (2.4) to $X \times \mathbb{P}^1 \times \mathbb{P}^1$ by p_{23} and using the commutativity of the above diagram, we get a short exact sequence on $X \times \mathbb{P}^1 \times \mathbb{P}^1$,

$$0 \longrightarrow (p_{12})^*q^*\mathcal{O}_{\mathbb{P}^1}(-1) \otimes (p_{13})^*q^*\mathcal{O}_{\mathbb{P}^1}(-1) \longrightarrow \mathcal{O}_{X \times \mathbb{P}^1 \times \mathbb{P}^1} \longrightarrow \mathcal{O}_{X \times \Delta_{\mathbb{P}^1}} \longrightarrow 0.$$

Tensoring this exact sequence with $(p_{12})^*E$ and then applying the functor $(p_{13})_*$, we have a long exact sequence

$$\begin{aligned} 0 &\longrightarrow (p_{13})_*[(p_{12})^*(E \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1))] \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \longrightarrow (p_{13})_*(p_{12})^*E \\ &\longrightarrow (p_{13})_*((p_{12})^*E|_{X \times \Delta_{\mathbb{P}^1}}) \longrightarrow R^1(p_{13})_*[(p_{12})^*(E \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1))] \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1). \end{aligned} \quad (2.5)$$

Consider the following commutative diagram,

$$\begin{array}{ccc} X \times \mathbb{P}^1 \times \mathbb{P}^1 & \xrightarrow{p_{13}} & X \times \mathbb{P}^1 \\ p_{12} \downarrow & & \downarrow p \\ X \times \mathbb{P}^1 & \xrightarrow{p} & X. \end{array}$$

Since all the maps in this diagram are flat, we have

$$R^1(p_{13})_*[(p_{12})^*(E \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1))] \cong p^*R^1p_*(E \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)) = 0.$$

It implies that the sequence

$$\begin{aligned} 0 &\longrightarrow (p_{13})_*[(p_{12})^*(E \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1))] \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \\ &\longrightarrow (p_{13})_*(p_{12})^*E \longrightarrow (p_{13})_*((p_{12})^*E|_{X \times \Delta_{\mathbb{P}^1}}) \longrightarrow 0 \end{aligned}$$

is exact.

We have $(p_{13})_*((p_{12})^*E|_{X \times \Delta_{\mathbb{P}^1}}) \cong E$. To complete the proof, we need to show that

$$(p_{13})_*[(p_{12})^*(E \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1))] \cong p^*E_0$$

and

$$(p_{13})_*(p_{12})^*E \cong p^*E_1$$

for some vector bundles E_0, E_1 on X . By Lemma 2.2.2, it is clear because

$$(p_{13})_*(p_{12}^*E)|_{F_p} \cong \mathcal{O}_{\mathbb{P}^1} \otimes H^0(\mathbb{P}^1, E|_{F_p})$$

and

$$(p_{13})_*[(p_{12})^*(E \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1))]|_{F_p} \cong \mathcal{O}_{\mathbb{P}^1} \otimes H^0(\mathbb{P}^1, E|_{F_p} \otimes \mathcal{O}_{\mathbb{P}^1}(-1))$$

for any F_p be any p -fiber. □

Let H be an ample divisor on $X \times \mathbb{P}^1$ and E be a vector bundle of rank r . We define the H -degree and H -slope of E by setting

$$\deg_H(E) = c_1(E).H \text{ and } \mu_H(E) = \frac{\deg_H(E)}{r}. \quad (2.6)$$

Then E is called H -(semi)stable if for any nonzero subsheaf $F \subseteq E$ we have $\mu_H(F) (\leq) \mu_H(E)$.

Theorem 2.2.4 (Bogomolov's inequality). *Let E be a vector bundle of rank r and H be an ample divisor on $X \times \mathbb{P}^1$. If E is H -semistable then*

$$\Delta(E) = (r-1)c_1^2(E) - 2rc_2(E) \leq 0,$$

where $c_1(E), c_2(E)$ are the first and second Chern classes of E .

Proof. Suppose that E is H -semistable but $\Delta(E) > 0$. Then we will construct a subsheaf $E' \subset E$ which destabilizes E .

I - **Construction of E'** . By Serre's vanishing theorem we can show that for any vector bundle E on $X \times \mathbb{P}^1$, there exists a line bundle L such that $R^1p_*(E \otimes L) = 0$. Since the semistability of E is unchanged when we tensor E with L and $\Delta(E) = \Delta(E \otimes L)$, by replacing E with $E \otimes L$ if necessary, we can assume that $R^1p_*(E \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)) = 0$. It follows from Proposition 2.2.3 that E has a resolution

$$0 \longrightarrow p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \longrightarrow p^*E_0 \longrightarrow E \longrightarrow 0, \quad (2.7)$$

where E_1, E_0 are vector bundles on X . Assume that E_i has rank r_i and degree d_i for $i = 1, 2$. By computing Chern characters up to numerical equivalence, we have

$$\text{ch}(p^*E_0) \equiv r_0 + d_0F_p, \text{ch}(p^*E_1) \equiv r_1 + d_1F_p, \text{ch}(q^*\mathcal{O}_{\mathbb{P}^1}(-1)) \equiv 1 - F_q,$$

$$\text{ch}(p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)) \equiv \text{ch}(p^*E_1) \text{ch}(q^*\mathcal{O}_{\mathbb{P}^1}(-1)) \equiv r_1 + d_1F_p - r_1F_q - d_1[pt],$$

$$\text{ch}(E) \equiv \text{ch}(p^*E_0) - \text{ch}(p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)) \equiv r_0 - r_1 + (d_0 - d_1)F_p + r_1F_q + d_1[pt].$$

Therefore

$$c_1(E) = \text{ch}_1(E) \equiv (d_0 - d_1)F_p + r_1F_q,$$

$$c_2(E) = \frac{c_1^2(E) - 2\text{ch}_2(E)}{2} \equiv (d_0r_1 - d_1r_1 - d_1)[pt].$$

It follows that

$$\Delta(E) = 2(r_0d_1 - r_1d_0) = 2r_0r_1(\mu(E_1) - \mu(E_0)).$$

Assume that $\Delta(E) > 0$, then $\mu(E_1) > \mu(E_0)$. Lemma 1.2.6 allows us to choose $\tau \in \mathbb{Q}$ such that

$$\frac{\text{rank}(E_{1(\tau)})}{r_1} > \frac{\text{rank}(E_{0(\tau)})}{r_0}.$$

We have $p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)|_{F_q} \subset p^*E_0|_{F_q}$ for any q -fiber $F_q \cong X$. Then $E_1 \subset E_0$ and hence $E_{1(\tau)} \subset E_{0(\tau)}$ by Corollary 1.2.5. Particularly, $E_{0(\tau)}$ and $E_{1(\tau)}$ are nontrivial. So

$$\frac{\text{rank}(E_{1(\tau)})}{r_1} > \frac{\text{rank}(E_{0(\tau)})}{r_0} \iff \frac{\text{rank}(E_{1(\tau)})}{\text{rank}(E_{0(\tau)})} > \frac{r_1}{r_0}.$$

We choose such τ that $\frac{\text{rank}(E_{1(\tau)})}{\text{rank}(E_{0(\tau)})}$ becomes maximal. Assume that $\text{rank}(E_{1(\tau)}) = r_{1\tau}$ and $\text{rank}(E_{0(\tau)}) = r_{0\tau}$. If $\mu(E_{1(\tau)}) > \mu(E_{0(\tau)})$, it follows from Lemma 1.2.6 that there exists $\rho \in \mathbb{Q}$ such that

$$\frac{\text{rank}((E_{1(\tau)})_{(\rho)})}{r_{1\tau}} > \frac{\text{rank}((E_{0(\tau)})_{(\rho)})}{r_{0\tau}}$$

where $(E_{1(\tau)})_{(\rho)}$ and $(E_{0(\tau)})_{(\rho)}$ are both nontrivial. It follows that

$$\frac{\text{rank}((E_{1(\tau)})_{(\rho)})}{\text{rank}((E_{0(\tau)})_{(\rho)})} > \frac{r_{1\tau}}{r_{0\tau}} > \frac{r_1}{r_0}.$$

This contradicts the choice of τ . Hence

$$\mu(E_{1(\tau)}) \leq \mu(E_{0(\tau)}). \quad (2.8)$$

Since $E_{1(\tau)}$ is subbundle of $E_{0(\tau)}$, the image of the composition

$$p^*E_{1(\tau)} \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \hookrightarrow p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \hookrightarrow p^*E_0$$

is contained in $p^*E_{0(\tau)}$. We obtain the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & p^*E_{1(\tau)} \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) & \longrightarrow & p^*E_{0(\tau)} & \longrightarrow & E'' \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \varphi \\ 0 & \longrightarrow & p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) & \longrightarrow & p^*E_0 & \longrightarrow & E \longrightarrow 0 \end{array}$$

Let $E' = \text{Im}(\varphi)$ and $K = \text{Ker}(\varphi)$. We have $\mathcal{O}_{\mathbb{P}^1}^{r_0\tau} \cong p^*E_{0(\tau)}|_{F_p} \rightarrow E''|_{F_p}$, equivalently, $E''|_{F_p}$ is globally generated. Since $K \subset p^*(E_1/E_{1(\tau)}) \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)$, $H^0(\mathbb{P}^1, K|_{F_p}) = 0$. Therefore $K \neq E''$ and then $E' \neq 0$.

II - E' **destabilizes** E . It follows from Proposition 2.1.4 that any ample divisor H on $X \times \mathbb{P}^1$ is in form $aF_p + bF_q$ for some $a, b \in \mathbb{Z}_{>0}$. Suppose that

$$c_1(E') \equiv m'F_p + n'F_q, c_1(E) = mF_p + nF_q,$$

then $\mu_H(E') = \frac{c_1(E') \cdot H}{r'} = \frac{m'b + n'a}{r'}$ and $\mu_H(E) = \frac{mb + na}{r}$. We have

$$\deg(E'|_{F_p}) = n', \deg(E'|_{F_q}) = m'.$$

If $\mu(E'|_{F_p}) > \mu(E|_{F_p})$ and $\mu(E'|_{F_q}) > \mu(E|_{F_q})$, then $m'/r' > m/r$ and $n'/r' > n/r$. It follows that

$$\mu_H(E') = \frac{an' + bm'}{r'} > \frac{am + bn}{r} = \mu_H(E).$$

Therefore, to show that E' destabilizes E , it is suffice to show that $E'|_{F_p}$ destabilizes $E|_{F_p}$ and $E'|_{F_q}$ destabilizes $E|_{F_q}$ for any F_p and F_q .

+ $\mu(E'|_{F_p}) > \mu(E|_{F_p})$: Let F_p be any p -fiber. Restricting the resolution (2.7) of E to F_p , we have an exact sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^1}^{r_1}(-1) \longrightarrow \mathcal{O}_{\mathbb{P}^1}^{r_0} \longrightarrow E|_{F_p} \longrightarrow 0.$$

It implies that

$$\text{rank}(E|_{F_p}) = r_0 - r_1, \deg(E|_{F_p}) = r_1.$$

Similarly,

$$\text{rank}(E''|_{F_p}) = r_{0\tau} - r_{1\tau}, \deg(E''|_{F_p}) = r_{1\tau}.$$

As we have seen, $\frac{r_{1\tau}}{r_{0\tau}} > \frac{r_1}{r_0}$, so

$$\mu(E''|_{F_p}) = \frac{r_{1\tau}}{r_{0\tau} - r_{1\tau}} > \mu(E|_{F_p}) = \frac{r_1}{r_0 - r_1}.$$

Restricting the following exact sequence

$$0 \longrightarrow K \longrightarrow E'' \longrightarrow E' \longrightarrow 0 \quad (2.9)$$

to F_p , we get an exact sequence

$$0 \longrightarrow K|_{F_p} \longrightarrow E''|_{F_p} \longrightarrow E'|_{F_p} \longrightarrow 0. \quad (2.10)$$

We have $\mu(E''|_{F_p}) = \frac{r_{1\tau}}{r_{0\tau} - r_{1\tau}} > 0$ by the choice of τ . Since

$$K \subset p^*(E_1/E_{1(\tau)}) \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)$$

then $K|_{F_p} \subset \mathcal{O}_{\mathbb{P}^1}^{r_1 - r_{1\tau}}$. It follows from Theorem 2.2.1 that

$$\mu(K|_{F_p}) \leq -1 < \mu(E''|_{F_p}).$$

From the exactness of (2.10),

$$\mu(K|_{F_p}) < \mu(E''|_{F_p}) \implies \mu(E'|_{F_p}) > \mu(E''|_{F_p}) > \mu(E|_{F_p}).$$

+ $\mu(E'|_{F_q}) > \mu(E|_{F_q})$: Let F_q be any q -fiber. Restricting (2.9) to F_q , we have an exact sequence

$$0 \longrightarrow K|_{F_q} \longrightarrow E''|_{F_q} \longrightarrow E'|_{F_q} \longrightarrow 0 \quad (2.11)$$

where $K|_{F_q} \subseteq E_1/E_{1(\tau)}$. By (iv) of Proposition 1.2.4, $\mu_{\max}(E_1/E_{1(\tau)}) < \tau$, hence

$$\mu(K|_{F_q}) \leq \mu_{\max}(E_1/E_{1(\tau)}) < \tau.$$

Consider the following exact sequence

$$0 \longrightarrow E_{1(\tau)} \longrightarrow E_{0(\tau)} \longrightarrow E''|_{F_q} \longrightarrow 0.$$

Since $\mu(E_{0(\tau)}) \geq \mu(E_{1(\tau)})$ (see (2.8)),

$$\mu(E''|_{F_q}) \geq \mu(E_{0(\tau)}) \geq \tau.$$

Hence $\mu(K|_{F_q}) < \mu(E''|_{F_q})$. It follows from the sequence (2.11) that

$$\mu(E'|_{F_q}) > \mu(E''|_{F_q}).$$

Similarly, since $\mu(E_1) > \mu(E_0)$ and the sequence

$$0 \longrightarrow E_1 \longrightarrow E_0 \longrightarrow E|_{F_q} \longrightarrow 0$$

is exact, we have

$$\mu(E_0) > \mu(E|_{F_q}).$$

It is clear by definition that $\mu(E_{0(\tau)}) \geq \mu(E_0)$. So we have

$$\mu(E'|_{F_q}) > \mu(E''|_{F_q}) \geq \mu(E_{0(\tau)}) \geq \mu(E_0) > \mu(E|_{F_q}).$$

□

Remark 2.2.5. Let E be a torsion free coherent sheaf on $X \times \mathbb{P}^1$ and $E^\vee = \text{Hom}_{\mathcal{O}_X}(E, \mathcal{O}_X)$ be the dual sheaf of E . Then $E^{\vee\vee} = (E^\vee)^\vee$ is a vector bundle of the same rank and $T := E^{\vee\vee}/E$ is a sheaf of rank zero. Consider the following exact sequence

$$0 \longrightarrow E \longrightarrow E^{\vee\vee} \longrightarrow T \longrightarrow 0$$

It is easy to see that $\Delta(E) = \Delta(E^{\vee\vee}) - 2 \text{rank}(E) \text{length}(T) \leq \Delta(E^{\vee\vee})$. Moreover, we have known that E is H -semistable if and only if $E^{\vee\vee}$ is H -semistable. Therefore, Bogomolov's inequality remains true for torsion free coherent sheaves.

2.3 Restriction theorems

Using Bogomolov's inequality, we can consider the semistability of vector bundles on $X \times \mathbb{P}^1$ restricted to a smooth curve.

Lemma 2.3.1 (cf. [13, Thm. 5.1]). *Let $0 = E_0 \subset E_1 \subset \dots \subset E_m = E$ be the Harder-Narasimhan filtration of E . Set $F_i = E_i/E_{i-1}$, $r_i = \text{rank}(F_i)$, $\mu_i = \mu(F_i)$. Then we have*

$$\frac{\Delta(E)}{r} = \sum_{i=1}^m \frac{\Delta(F_i)}{r_i} + \frac{1}{r} \sum_{i < j} r_i r_j \left(\frac{c_1(F_i)}{r_i} - \frac{c_1(F_j)}{r_j} \right)^2. \quad (2.12)$$

Proof. We have

$$\begin{aligned} \Delta(E) &= (r-1)c_1^2(E) - 2rc_2(E) \\ &= r(c_1^2(E) - 2c_2(E)) - c_1^2(E) \\ &= 2 \text{ch}_0(E) \text{ch}_2(E) - \text{ch}_1^2(E), \end{aligned}$$

where $\text{ch}_i(E)$ is the i^{th} term in the expression of the Chern character of E :

$$\text{ch}(E) = r + c_1(E) + \frac{(c_1^2(E) - 2c_2(E))}{2}.$$

Since Chern character is additive with respect to an exact sequence, we have

$$\frac{\Delta(E)}{r} = 2 \text{ch}_2(E) - \frac{c_1^2(E)}{r} = 2 \sum_{i=1}^m \text{ch}_2(F_i) - \frac{(\sum_{i=1}^m c_1(F_i))^2}{r},$$

$$\sum_{i=1}^m \frac{\Delta(F_i)}{r_i} = 2 \sum_{i=1}^m \text{ch}_2(F_i) - \sum_{i=1}^m \frac{c_1^2(F_i)}{r_i}.$$

Therefore the quality (2.12) is equivalent to

$$\sum_{i=1}^m \frac{c_1^2(F_i)}{r_i} = \frac{1}{r} \left[\left(\sum_{i=1}^m c_1(F_i) \right)^2 + \sum_{i < j} r_i r_j \left(\frac{c_1(F_i)}{r_i} - \frac{c_1(F_j)}{r_j} \right)^2 \right]. \quad (2.13)$$

But it is clear because

$$\begin{aligned} \left(\sum_{i=1}^m c_1(F_i) \right)^2 + \sum_{i < j} r_i r_j \left(\frac{c_1(F_i)}{r_i} - \frac{c_1(F_j)}{r_j} \right)^2 &= \sum_{i=1}^m c_1^2(F_i) + \sum_{i < j} \left(\frac{r_j c_1^2(F_i)}{r_i} + \frac{r_i c_1^2(F_j)}{r_j} \right) \\ &= \sum_{j=1}^m c_1^2(F_j) + \sum_{j=1}^m \frac{(\sum_{i \neq j} r_i) c_1^2(F_j)}{r_j} \\ &= r \sum_{j=1}^m \frac{c_1^2(F_j)}{r_j}. \end{aligned}$$

□

Proposition 2.3.2 (cf. [13, Thm. 5.1]). *Let E be any rank r torsion free sheaf on $X \times \mathbb{P}^1$ and H be an ample divisor. Then we have*

$$H^2 \cdot \Delta(E) \leq r^2 (\mu_{\max} - \mu)(\mu - \mu_{\min}).$$

Proof. Let $0 = E_0 \subset E_1 \subset \dots \subset E_m = E$ be the Harder-Narasimhan filtration of E . Set $F_i = E_i/E_{i-1}$, $r_i = \text{rank}(F_i)$, $\mu_i = \mu(F_i)$. We have

$$\frac{\Delta(E)}{r} = \sum \frac{\Delta(F_i)}{r_i} + \frac{1}{r} \sum_{i < j} r_i r_j \left(\frac{c_1(F_i)}{r_i} - \frac{c_1(F_j)}{r_j} \right)^2 \leq \frac{1}{r} \sum_{i < j} r_i r_j \left(\frac{c_1(F_i)}{r_i} - \frac{c_1(F_j)}{r_j} \right)^2$$

since $\Delta(F_i) \leq 0$ for all i . It follows from (2.2) that

$$H^2 \left(\frac{c_1(F_i)}{r_i} - \frac{c_1(F_j)}{r_j} \right)^2 \leq \left(\left(\frac{c_1(F_i)}{r_i} - \frac{c_1(F_j)}{r_j} \right) H \right)^2.$$

Since H is ample, $H^2 > 0$. We have

$$\begin{aligned} \frac{\Delta(E)}{r} &\leq \frac{1}{r H^2} \sum_{i < j} r_i r_j \left(\left(\frac{c_1(F_i)}{r_i} - \frac{c_1(F_j)}{r_j} \right) H \right)^2 \\ &= \frac{1}{r H^2} \sum_{i < j} r_i r_j (\mu_i - \mu_j)^2. \end{aligned}$$

By the definition of the Harder-Narasimhan filtration, we have

$$\mu_{\max} = \mu_1 > \mu_2 > \dots > \mu_m = \mu_{\min},$$

$$r = \sum_{i=1}^m r_i,$$

$$r\mu(E) = \sum_{i=1}^m r_i\mu_i.$$

Moreover

$$\begin{aligned} \sum_{i < j} r_i r_j (\mu_i - \mu_j)^2 &= r \left(\sum_{i=1}^{m-1} \left(\sum_{j \leq i} r_j (\mu_j - \mu) \right) (\mu_i - \mu_{i+1}) \right) \\ &\leq r \left(\sum_{i=1}^{m-1} \left(\sum_{j \leq i} r_j (\mu_1 - \mu) \right) (\mu_i - \mu_{i+1}) \right) \\ &= r(\mu_1 - \mu) \left(\sum_{i=1}^{m-1} \left(\sum_{j \leq i} r_j \right) (\mu_i - \mu_{i+1}) \right) \\ &= r(\mu_{\max} - \mu) \left(\sum_{i=1}^m r_i \mu_i - r\mu_m \right) \\ &= r^2(\mu_{\max} - \mu)(\mu - \mu_{\min}). \end{aligned}$$

So we have

$$\frac{\Delta(E)}{r} \leq \frac{1}{rH^2} \sum_{i < j} r_i r_j (\mu_i - \mu_j)^2 \leq \frac{r^2(\mu_{\max} - \mu)(\mu - \mu_{\min})}{rH^2}.$$

It implies that

$$H^2\Delta(E) \leq r^2(\mu_{\max} - \mu)(\mu - \mu_{\min}).$$

□

Theorem 2.3.3 ([13, Thm. 5.2]). *Let E be a vector bundle on $X \times \mathbb{P}^1$ of rank $r \geq 2$. Assume that E is H -stable. Let $C \in |mH|$ be a smooth curve. If $m \geq \frac{1-r}{r}\Delta(E) + 1$ then $E|_C$ is stable.*

Proof. Assume that $E|_C$ is not stable. Then there exists a quotient Q of $E|_C$ which is a vector bundle of rank R and degree d on C such that $\mu(E|_C) \geq \mu(Q)$. Let E' be the kernel of the composition $E \rightarrow E|_C \rightarrow Q$ and consider the following exact sequence

$$0 \rightarrow E' \rightarrow E \rightarrow Q \rightarrow 0,$$

where Q is considered as a sheaf of rank zero on $X \times \mathbb{P}^1$. By computing with Chern character, we have

$$\text{ch}(Q) \equiv RC + \left(d - \frac{RC^2}{2}\right)[pt].$$

Therefore,

$$\text{ch}(E') = \text{ch}(E) - \text{ch}(Q) \equiv r + (c_1(E) - RmH) + \left(\frac{\Delta(E) + c_1^2(E)}{2r} - d + \frac{Rm^2H^2}{2}\right)[pt].$$

It follows that

$$\begin{aligned}\Delta(E') &= 2\operatorname{ch}_0(E')\operatorname{ch}_2(E') - \operatorname{ch}_1^2(E') \\ &= \Delta(E) + 2rR(\mu(E|_C) - \mu(Q)) + m^2H^2R(r - R) \\ &\geq \Delta(E) + m^2H^2R(r - R).\end{aligned}$$

If E' is semistable, $\Delta(E') \leq 0$ by Bogomolov's inequality. So we have

$$-\Delta(E) \geq m^2H^2R(r - R) \geq (r - 1)m^2H^2$$

since $1 \leq R \leq r - 1$. It implies then

$$\frac{1 - r}{r}\Delta(E) + 1 \geq \frac{(r - 1)^2}{r}m^2H^2 + 1.$$

But it is a contradiction since $r \geq 2$, $H^2 \geq 1$ and $m \geq \frac{1-r}{r}\Delta(E) + 1$. So E' can not be semistable. In that case, we have

$$\mu_{\max}(E') - \mu(E') = \mu_{\max}(E') - \mu(E) + \frac{R}{r}mH^2 = \frac{R}{r}mH^2 - \frac{dr' - d'r}{rr'},$$

where r', d' are the rank and degree of the maximal destabilizing subbundle of E' . Since E is stable, $dr' - d'r \geq 1$. Moreover $1 \leq r' \leq r - 1$, then

$$\mu_{\max}(E') - \mu(E') \leq \frac{R}{r}mH^2 - \frac{1}{r(r - 1)}.$$

Consider the following diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & E' & \longrightarrow & S & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & E(-C) & \longrightarrow & E & \longrightarrow & E|_C \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & Q & \xrightarrow{\operatorname{Id}} & Q & & \\ & & \downarrow & & \downarrow & & \\ & & 0 & & 0 & & \end{array}$$

where S is the kernel of $E|_C \rightarrow Q$ on C and the rows are exact. By the kernel-cokernel exact sequence,

$$0 \longrightarrow E(-C) \longrightarrow E' \longrightarrow S \longrightarrow 0$$

is an exact sequence where S is a sheaf considered on $X \times \mathbb{P}^1$. In particular $E'^\vee \subset E(-C)^\vee$ and $E(-C)^\vee$ are still stable. Therefore

$$\mu(E') - \mu_{\min}(E') = \mu(E(-C)) + \frac{r - R}{r}mH^2 - \mu_{\min}(E')$$

$$\begin{aligned}
 &= \mu_{\max}(E'^{\vee}) - \mu(E(-C)^{\vee}) + \frac{r-R}{r}mH^2 \\
 &\leq \frac{r-R}{r}mH^2 - \frac{1}{r(r-1)}.
 \end{aligned}$$

Now we apply Proposition 2.3.2 for E' and obtain

$$\begin{aligned}
 0 &\geq H^2\Delta(E') - r^2(\mu_{\max}(E') - \mu(E'))(\mu(E') - \mu_{\min}(E')) \\
 &\geq H^2\Delta(E) + m^2H^4R(r-R) - r^2\left(\frac{R}{r}mH^2 - \frac{1}{r(r-1)}\right)\left(\frac{r-R}{r}mH^2 - \frac{1}{r(r-1)}\right) \\
 &= H^2\Delta(E) + \frac{rH^2}{r-1}m + \frac{1}{(r-1)^2}.
 \end{aligned}$$

Therefore

$$\frac{rH^2}{1-r}m \geq H^2\Delta(E) + \frac{1}{(r-1)^2} \iff m \leq \frac{1-r}{r}\Delta(E) - \frac{1}{r(r-1)H^2}.$$

This contradicts the assumption that $m \geq \frac{1-r}{r}\Delta(E) + 1$. \square

Corollary 2.3.4 (cf. [13, Cor. 5.4]). *Let E be a vector bundle of rank $r \geq 2$ on $X \times \mathbb{P}^1$ and $C \in |mH|$ be a general smooth projective curve. Assume that E is H -semistable. If*

$$m \geq \frac{1-r}{r}\Delta(E) + 1$$

then $E|_C$ is semistable.

Proof. Assume that we have a short exact sequence of vector bundles on $X \times \mathbb{P}^1$

$$0 \longrightarrow E_1 \longrightarrow E \longrightarrow E_2 \longrightarrow 0,$$

where $\mu_H(E_i) = \mu_H(E)$. Let $C \in |mH|$ be a general smooth projective curve. By restricting to C , the sequence

$$0 \longrightarrow E_1|_C \longrightarrow E|_C \longrightarrow E_2|_C \longrightarrow 0$$

is still exact and $\mu(E_1|_C) = \mu(E_2|_C) = \mu(E|_C)$. It is easy to see that $E|_C$ is semistable if $E_i|_C$ is semistable for $i = 1, 2$. By induction on the length of a Jordan-Hölder filtration which depends only on E itself, it is enough to assume that E has a Jordan-Hölder filtration of length 2,

$$0 = E_0 \subset E_1 \subset E.$$

Set $E_2 = E/E_1$ and consider the exact sequence

$$0 \longrightarrow E_1|_C \longrightarrow E|_C \longrightarrow E_2|_C \longrightarrow 0$$

where $\mu(E_1|_C) = \mu(E_2|_C) = \mu(E|_C)$. By Theorem 2.3.3, $E_i|_C$ are semistable if

$$m \geq \frac{1-r_i}{r_i}\Delta(E_i) + 1.$$

As in (2.12), we have

$$\frac{\Delta(E)}{r} = \frac{\Delta(E_1)}{r_1} + \frac{\Delta(E_2)}{r_2} + \frac{r_1 r_2}{r} \left(\frac{c_1(E_1)}{r_1} - \frac{c_1(E_2)}{r_2} \right)^2.$$

It follows that

$$\begin{aligned} \frac{1-r}{r} \Delta(E) + 1 &= \frac{1-r_1}{r_1} \Delta(E_1) + 1 + \frac{1-r_2}{r_2} \Delta(E_2) \\ &\quad + \left[(1-r) \frac{r_1 r_2}{r} \left(\frac{c_1(E_1)}{r_1} - \frac{c_1(E_2)}{r_2} \right)^2 - \frac{r_2}{r_1} \Delta(E_1) - \frac{r_1}{r_2} \Delta(E_2) \right] \end{aligned}$$

We have $\Delta(E_i) \leq 0$ since E_i are semistable. Moreover $\mu(E_2) = \mu(E_1)$. It follows from (2.1) that

$$\left(\frac{c_1(E_1)}{r_1} - \frac{c_1(E_2)}{r_2} \right)^2 \leq 0.$$

So we have

$$\begin{aligned} \frac{1-r}{r} \Delta(E) + 1 &\geq \frac{1-r_1}{r_1} \Delta(E_1) + 1 + \frac{1-r_2}{r_2} \Delta(E_2) \\ &\geq \max \left\{ \frac{1-r_1}{r_1} \Delta(E_1) + 1, \frac{1-r_2}{r_2} \Delta(E_2) + 1 \right\}. \end{aligned}$$

Hence for any

$$m \geq \frac{1-r}{r} \Delta(E) + 1,$$

$E_i|_C$ is semistable. Therefore, $E|_C$ is semistable.

□

Chapter 3

Triples and their moduli spaces

3.1 Triples and their stability

Let X be a smooth projective curve over an algebraically closed field k . We consider the first notions of triples and their stability.

Definition 3.1.1. A triple T on X consists of two vector bundles E_1, E_2 on X , together with a morphism $\varphi : E_2 \longrightarrow E_1$ and it is denoted by $T = (E_2 \xrightarrow{\varphi} E_1)$ or simply $T = (E_2, E_1, \varphi)$.

Let $T = (E_2, E_1, \varphi)$ and $T' = (E'_2, E'_1, \varphi')$ be triples on X . Then a morphism of triples, $f : T \longrightarrow T'$, is a pair (f_2, f_1) of morphisms such that the following diagram commutes

$$\begin{array}{ccc} E_2 & \xrightarrow{\varphi} & E_1 \\ \downarrow f_2 & & \downarrow f_1 \\ E'_2 & \xrightarrow{\varphi'} & E'_1. \end{array} \quad (3.1)$$

In particular, f is an isomorphism if f_1 and f_2 are both isomorphisms.

If $\text{rank}(E_i) = r_i$ and $\deg(E_i) = d_i$ for $i = 1, 2$, then we say that T is of type (r_1, r_2, d_1, d_2) .

To introduce the notion of (semi)stability for triples, we first precise the subtriples of a given one and then define the slope which depends on a real parameter.

Definition 3.1.2. A triple $T' = (E'_2, E'_1, \varphi')$ is called a subtriple of T if E'_i is a subbundle of E_i for $i = 1, 2$ and the commutative diagram (3.1) holds for $f = (j_2, j_1)$ where $j_i : E'_i \hookrightarrow E_i$ are the inclusions. If $E'_1 = E'_2 = 0$ then T' is called trivial subtriple. T' is called proper subtriple if it is nontrivial and $T' \neq T$.

Example 3.1.3. Let $T = (L, E, \varphi)$ be a triple where L is a line bundle on X . Then any subtriple T' of T is either $T' = (0, E', 0)$ where $E' \subset E$ is any subbundle or $T' = (L, E', \varphi)$ where E' is a subbundle of E such that $\varphi(L) \subset E'$.

Definition 3.1.4. Let $T = (E_2, E_1, \varphi)$ be a triple of type (r_1, r_2, d_1, d_2) . For any real number $\alpha \in \mathbb{R}$, we define

$$\deg_\alpha(T) = d_1 + d_2 - 2\alpha r_1$$

and

$$\mu_\alpha(T) = \frac{\deg_\alpha(T)}{r_1 + r_2} = \frac{d_1 + d_2 - 2\alpha r_1}{r_1 + r_2}. \quad (3.2)$$

The triple T is called α -(semi)stable if for all nontrivial subtriples T' of T , $\mu_\alpha(T')(\leq)\mu_\alpha(T)$.

Example 3.1.5. We consider the simplest triples $T = (0, E, 0)$ or $T = (E, 0, 0)$. Then

$$\mu_\alpha(T')(\leq)\mu_\alpha(T) \iff \mu(E')(\leq)\mu(E)$$

where $E' \subsetneq E$ and $T' = (0, E', 0)$ or $T' = (E', 0, 0)$. Hence the α -(semi)stability of T is in fact equivalent to the (semi)stability of E for any α . From now on, we just consider triples T with E_1 and E_2 are both nontrivial.

Remark 3.1.6. The above definition of α -degree and α -slope slightly differ from the definitions given in [2],[4], etc, where the authors defined $\deg_\alpha(T) = d_1 + d_2 + \alpha r_2$. This modification does not change the (semi)stability of triples and is useful when we connect the semistability of extension bundles on $X \times \mathbb{P}^1$ defined by T with the semistability of T itself.

Lemma 3.1.7. For a short exact sequence of triples on X ,

$$\begin{array}{ccccccc} 0 & \longrightarrow & E'_2 & \longrightarrow & E_2 & \longrightarrow & E''_2 \longrightarrow 0 \\ & & \downarrow \varphi' & & \downarrow \varphi & & \downarrow \varphi'' \\ 0 & \longrightarrow & E'_1 & \longrightarrow & E_1 & \longrightarrow & E''_1 \longrightarrow 0, \end{array} \quad (3.3)$$

we have

$$\mu_\alpha((E'_2, E'_1, \varphi'))(\leq)\mu_\alpha((E_2, E_1, \varphi)) \iff \mu_\alpha((E_2, E_1, \varphi))(\leq)\mu_\alpha((E''_2, E''_1, \varphi')).$$

Proof. Since the rows are exact, we have

$$d_i = d'_i + d''_i, r_i = r'_i + r''_i$$

for $i = 1, 2$. It follows that

$$\mu_\alpha((E_2, E_1, \varphi)) = \frac{\deg_\alpha(E_2, E_1, \varphi)}{r_1 + r_2} = \frac{\deg_\alpha(E'_2, E'_1, \varphi') + \deg_\alpha(E''_2, E''_1, \varphi')}{(r'_1 + r'_2) + (r''_1 + r''_2)}.$$

Hence

$$\frac{\deg_\alpha(E_2, E_1, \varphi)}{r_1 + r_2}(\geq)\frac{\deg_\alpha(E'_2, E'_1, \varphi')}{r'_1 + r'_2} \iff \frac{\deg_\alpha(E_2, E_1, \varphi)}{r_1 + r_2}(\leq)\frac{\deg_\alpha(E''_2, E''_1, \varphi')}{r''_1 + r''_2}.$$

□

Let $T = (E_2, E_1, \varphi)$ be a triple and L be a line bundle on X . We define the triple $T \otimes L$ to be the triple $(E_2 \otimes L, E_1 \otimes L, \varphi \otimes \text{Id}_L)$.

Lemma 3.1.8. *Let $T = (E_2, E_1, \varphi)$ be a triple on X and L a line bundle. Then T is α -semistable if and only if $T \otimes L$ is α -semistable.*

Proof. We have

$$\mu_\alpha(T \otimes L) = \mu_\alpha(T) + \deg(L).$$

Moreover, any subtriple of $T \otimes L = (E_2 \otimes L, E_1 \otimes L, \varphi \otimes \text{Id}_L)$ is induced from a subtriple of T by tensoring with L . Hence T is α -semistable if and only if $T \otimes L$ is. \square

It is well known that any stable vector bundle E is simple, i.e. $\text{End } E \cong k$. For stable triples, an analogous result holds true.

Definition 3.1.9. *Let $T = (E_2, E_1, \varphi)$ be a triple. Let us set*

$$\text{End}(T) = \{(u, v) \in \text{End}(E_2) \oplus \text{End}(E_1) \mid \varphi u = v\varphi\}.$$

We say that T is simple if $\text{End}(T) \cong k$, i.e. the only elements in $\text{End}(T)$ are of the form $(\lambda \text{Id}_{E_2}, \lambda \text{Id}_{E_1})$ for some $\lambda \in k$.

Proposition 3.1.10 ([2, Prop. 3.10]). *Let $T = (E_2, E_1, \varphi)$ be a α -stable triple and $(u, v) \in \text{End}(T)$. Then either (u, v) is trivial or both u, v are isomorphisms.*

Proof. Suppose that u and v are both neither nontrivial nor isomorphism. Then

$$T' = (\text{Ker } u, \text{Ker } v, \varphi) \text{ and } T'' = (\text{Im } u, \text{Im } v, \varphi)$$

are nontrivial subtriples of T . T is α -stable, then

$$\mu_\alpha(T') < \mu_\alpha(T) \text{ and } \mu_\alpha(T'') < \mu_\alpha(T). \quad (3.4)$$

We have an exact sequence of triples

$$0 \longrightarrow T' \longrightarrow T \longrightarrow T'' \longrightarrow 0.$$

It follows from lemma 3.1.7 that $\mu_\alpha(T'') > \mu_\alpha(T)$ if $\mu_\alpha(T) < \mu_\alpha(T')$. This contradicts (3.4). \square

Corollary 3.1.11 ([2, Cor. 3.12]). *If T is α -stable, then it is simple.*

For any triple $T = (E_2, E_1, \varphi)$ there is always a dual triple $T^\vee = (E_1^\vee, E_2^\vee, \varphi^\vee)$, where φ^\vee is the transpose of φ , i.e. the image of φ via the canonical isomorphism

$$\text{Hom}(E_2, E_1) \cong \text{Hom}(E_1^\vee, E_2^\vee).$$

From the one-to-one correspondence between subbundles of a given bundle E and subbundles of E^\vee , it follows that there is also a one-to-one correspondence between subtriples of T and subtriples of T^\vee . Therefore the α -(semi)stability of T is related to that of T^\vee .

Proposition 3.1.12 ([2, Prop. 3.16]). *The triple T is α -(semi)stable if and only if its dual triple T^\vee is α -(semi)stable*

Proof. By symmetric property, it is enough to show that T is α -(semi)stable if T^\vee is. Let $T' = (E'_2, E'_1, \varphi')$ be nontrivial subtriple of T and T'' be the quotient of T by T' . Then we have a short exact sequence of triples,

$$0 \longrightarrow (T'')^\vee \longrightarrow T^\vee \longrightarrow (T')^\vee \longrightarrow 0.$$

As T^\vee is α -(semi)stable, we have $\mu_\alpha(T^\vee)(\geq)\mu_\alpha((T'')^\vee)$. It follows from Lemma 3.1.7 that $\mu_\alpha(T^\vee)(\leq)\mu_\alpha((T')^\vee)$. But we have

$$\begin{aligned} \mu_\alpha(T^\vee)(\leq)\mu_\alpha((T')^\vee) &\iff \frac{d_1 + d_2 + 2\alpha r_2}{r_1 + r_2}(\geq)\frac{d'_1 + d'_2 + 2\alpha r'_2}{r'_1 + r'_2} \\ &\iff \frac{d_1 + d_2 - 2\alpha r_1 + 2\alpha(r_1 + r_2)}{r_1 + r_2}(\geq)\frac{d'_1 + d'_2 - 2\alpha r'_1 + 2\alpha(r'_1 + r'_2)}{r'_1 + r'_2} \\ &\iff \mu_\alpha(T)(\geq)\mu_\alpha(T'). \end{aligned}$$

Hence T is α -(semi)stable. □

3.2 Constraints on the parameter α

The constraints on α for α -(semi)stability of triples were first given by Bradlow and García-Prada in [2] where the authors defined the α -degree of $T = (E_2, E_1, \varphi)$ to be $d_1 + d_2 + \alpha r_2$. Using our definition of the α -degree

$$\deg_\alpha(T) := d_1 + d_2 - 2\alpha r_1,$$

we have,

$$\mu_\alpha(T) = \frac{\deg_\alpha(T)}{r_1 + r_2} = \frac{d_1 + d_2 - 2\alpha r_1}{r_1 + r_2} = \frac{d_1 + d_2 + 2\alpha r_2}{r_1 + r_2} - 2\alpha.$$

It follows that T is α -(semi)stable if and only if T is 2α -(semi)stable by definition of Bradlow and García-Prada. Hence the constraints on α with respect to our definition can be induced by [2, Sec 3.3].

Proposition 3.2.1 (cf. [2, Lm. 3.5]). *Let $T = (E_2, E_1, 0)$ be a triple. Then T is α -semistable if and only if $\alpha = \frac{\mu(E_1) - \mu(E_2)}{2}$ and both bundles E_1, E_2 are semistable. In this case T can not be α -stable.*

Proposition 3.2.2 (cf. [2, Prop. 3.17]). *Let $T = (E_2, E_1, \varphi)$ be a α -(semi)stable triple where $\varphi \neq 0$. Then $\alpha(\geq)\frac{\mu(E_1) - \mu(E_2)}{2}$ and $\alpha(\geq)0$.*

Proposition 3.2.3 (cf. [2, Prop. 3.18]). *Let $T = (E_2, E_1, \varphi)$ be a triple with $r_1 \neq r_2$ and $\varphi \neq 0$. If T is α -(semi)stable then*

$$\alpha(\leq)(1 + \frac{r_1 + r_2}{|r_1 - r_2|}) \frac{\mu(E_1) - \mu(E_2)}{2}. \quad (3.5)$$

Proof. Let $I = \text{Im } \varphi$ and $K = \text{Ker } \varphi$. Assume that $r_1 > r_2$ then φ can not be surjective. Then $T'' = (E_2, I, \varphi)$ is proper subtriple of T .

If φ is not injective, $T' = (K, 0, \varphi)$ is also nontrivial. As T is α -(semi)stable, we have

$$\mu_\alpha(T')(\leq)\mu_\alpha(T) \iff r_K(d_1 + d_2) - d_K(r_1 + r_2)(\geq)2\alpha r_1 r_K \quad (3.6)$$

and

$$\begin{aligned} \mu_\alpha(T'')(\leq)\mu_\alpha(T) &\iff \frac{2d_2 - d_K - 2\alpha(r_2 - r_K)}{2r_2 - r_K}(\leq)\frac{d_1 + d_2 - 2\alpha r_1}{r_1 + r_2} \\ &\iff 2(d_1 r_2 - d_2 r_1) + d_K(r_1 + r_2) - r_K(r_1 + r_2)(\geq)2\alpha(r_2(r_1 - r_2) + r_K(r_1 + r_2)) - 2\alpha_1 r_K. \end{aligned} \quad (3.7)$$

Let (3.6) add to (3.7), we get

$$d_1 r_2 - d_2 r_1(\geq)\alpha(r_2(r_1 - r_2) + r_K(r_1 + r_2)).$$

As $\varphi \neq 0$ and T is α -(semi)stable, we have $\alpha(\geq)0$. Therefore

$$d_1 r_2 - d_2 r_1(\geq)\alpha r_2(r_1 - r_2) \quad (3.8)$$

since $r_K > 0$.

If φ is injective, i.e. $K = 0$ then

$$\mu_\alpha(T'')(\leq)\mu_\alpha(T) \iff d_1 r_2 - d_2 r_1(\geq)\alpha r_2(r_1 - r_2).$$

So we always have

$$d_1 r_2 - d_2 r_1(\geq)\alpha(r_2(r_1 - r_2)).$$

It implies that

$$\alpha(\leq)(1 + \frac{r_1 + r_2}{|r_1 - r_2|}) \frac{(\mu(E_1) - \mu(E_2))}{2}.$$

If $r_1 < r_2$ then the dual triple $T^\vee = (E_1^\vee, E_2^\vee, \varphi^\vee)$ is also α -(semi)stable by Propostion 3.1.12. Applying the same computation for T^\vee , we get

$$\begin{aligned} \deg(E_2^\vee) \text{rank}(E_1^\vee) - \deg(E_1^\vee) \text{rank}(E_2^\vee)(\geq)\alpha \text{rank}(E_1^\vee)(\text{rank}(E_2^\vee) - \text{rank}(E_1^\vee)) \\ \iff d_1 r_2 - d_2 r_1(\geq)\alpha r_1(r_2 - r_1) \\ \iff \alpha(\leq)(1 + \frac{r_1 + r_2}{|r_1 - r_2|}) \frac{\mu(E_1) - \mu(E_2)}{2}. \end{aligned}$$

□

If $r_1 = r_2$ then T could be α -(semi)stable for any $\alpha \geq \frac{\mu(E_1) - \mu(E_2)}{2}$. For example, the triple $T = (\mathcal{O}_X, \mathcal{O}_X, \text{Id}_{\mathcal{O}_X})$, then for any $\alpha(\geq)0$, T is α -(semi)stable.

3.3 The corresponding extensions on $X \times \mathbb{P}^1$ of triples

3.3.1 The construction of extensions

Let $T = (E_2 \xrightarrow{\varphi} E_1)$ be a triple on X and $\alpha = \frac{a}{b} \in \mathbb{Q}$ such that $\gcd(a, b) = 1$. We will construct a vector bundle E_T on $X \times \mathbb{P}^1$ such that its semistability with respect to an ample divisor $H(\alpha)$ is equivalent to the α -semistability of T .

On \mathbb{P}^1 we have the Euler sequence

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^1}(-2) \longrightarrow H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1)) \otimes \mathcal{O}_{\mathbb{P}^1}(-1) \xrightarrow{e} \mathcal{O}_{\mathbb{P}^1} \longrightarrow 0. \quad (3.9)$$

Let $p : X \times \mathbb{P}^1 \longrightarrow X$ and $q : X \times \mathbb{P}^1 \longrightarrow \mathbb{P}^1$ be the projections. By pulling back the exact sequence (3.9) by q and then taking tensor product with p^*E_1 , we obtain a short exact sequence

$$0 \longrightarrow p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2) \longrightarrow p^*E_1 \otimes V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \xrightarrow{\psi} p^*E_1 \longrightarrow 0 \quad (3.10)$$

where $V := H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$ and $\psi := \text{Id}_{p^*E_1} \otimes q^*e$.

Let us set E_T as the pullback of two morphisms

$$\psi : p^*E_1 \otimes V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \longrightarrow p^*E_1$$

and

$$p^*\varphi : p^*E_2 \longrightarrow p^*E_1.$$

It is locally given by

$$E_T(U) = \{(\eta, \nu) \in p^*E_1 \otimes V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)(U) \oplus p^*E_2(U) \mid (p^*\varphi)(\nu) = \psi(\eta)\}$$

for any open subset $U \subseteq X \times \mathbb{P}^1$. Since ψ is surjective, we get also a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2) & \longrightarrow & E_T & \longrightarrow & p^*E_2 \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2) & \longrightarrow & p^*E_1 \otimes V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) & \longrightarrow & p^*E_1 \longrightarrow 0 \end{array} \quad (3.11)$$

with exact rows. Furthermore, we have an exact sequence

$$0 \longrightarrow E_T \longrightarrow p^*E_1 \otimes V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \oplus p^*E_2 \longrightarrow p^*E_1 \longrightarrow 0. \quad (3.12)$$

Lemma 3.3.1. *Let $T = (E_2 \xrightarrow{\varphi} E_1)$, $T' = (E'_2 \xrightarrow{\varphi'} E'_1)$ be triples on X . Then any morphism $f : T \longrightarrow T'$ induces a morphism $\bar{f} : E_T \longrightarrow E_{T'}$ of vector bundles on $X \times \mathbb{P}^1$ defined by T and T' , respectively. In particular, if f is injective then \bar{f} is.*

Proof. Assume that $f = (f_2, f_1) : T \longrightarrow T'$, i.e. we have a commutative diagram

$$\begin{array}{ccc} E_2 & \xrightarrow{\varphi} & E_1 \\ \downarrow f_2 & & \downarrow f_1 \\ E'_2 & \xrightarrow{\varphi'} & E'_1. \end{array}$$

By the construction of E_T and $E_{T'}$, we have the following diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & E_{T'} & \longrightarrow & p^*E'_1 \otimes V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \oplus p^*E'_2 & \longrightarrow & p^*E'_2 \longrightarrow 0 \\ & & & & \downarrow & & \downarrow \\ 0 & \longrightarrow & E_T & \longrightarrow & p^*E_1 \otimes V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \oplus p^*E_2 & \longrightarrow & p^*E_2 \longrightarrow 0 \end{array}$$

where the square on the right hand side commutes. From the exactness of the rows, there exists a morphism $\bar{f} : E_{T'} \longrightarrow E_T$ making the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & E_{T'} & \longrightarrow & p^*E'_1 \otimes V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \oplus p^*E'_2 & \longrightarrow & p^*E'_2 \longrightarrow 0 \\ & & \downarrow \bar{f} & & \downarrow & & \downarrow \\ 0 & \longrightarrow & E_T & \longrightarrow & p^*E_1 \otimes V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \oplus p^*E_2 & \longrightarrow & p^*E_2 \longrightarrow 0 \end{array}$$

commutative. It is clear that, \bar{f} is injective if f is injective. \square

3.3.2 $\mathrm{SL}(2)$ -invariant subbundles

It follows from Lemma 3.3.1 that any subtriple of T defines a subbundle of E_T . But it is obviously not true that every subbundle of E_T is obtained by this way. Hence we would like to look for the subbundles of E_T which are defined by subtriples of T .

The graded ring $S = k[x, y]$ is generated by S_1 , the space of homogeneous polynomials of degree 1, as k -algebra. On $\mathbb{P}^1 = \mathrm{Proj} S$, any $s \in S_d$, a homogeneous polynomial of degree d , determines in a natural way a global section of $\mathcal{O}_{\mathbb{P}^1}(d)$. Then x, y determine the standard basis of k -vector space $V = H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$. On V , $\mathrm{SL}(2) := \mathrm{SL}_2(V)$ acts naturally by

$$g(Ax + By) = \begin{pmatrix} A \\ B \end{pmatrix}^t g \begin{pmatrix} x \\ y \end{pmatrix}$$

for any $g \in \mathrm{SL}(2)$ and $Ax + By \in V$. We then have a natural action of $\mathrm{SL}(2)$ on S_d given by

$$g(s(x, y)) = s(g(x), g(y))$$

for any $s = s(x, y) \in S_d$ and $g \in \mathrm{SL}(2)$. This action is compatible with the action of $\mathrm{SL}(2)$ on \mathbb{P}^1 given by

$$g \circ P = (du - bv : -cu + av)$$

for any $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2)$ and $P = (u : v) \in \mathbb{P}^1$. Assume that $P = Z(\mathfrak{p}) \in \mathbb{P}^1$, we have

$$\mathcal{O}_{\mathbb{P}^1}(n)_P \cong S(n)_{(\mathfrak{p})}$$

where $S(n)_i = S_{n+i}$ for any $i \geq 0, n \in \mathbb{Z}$ and an isomorphism

$$S(n)_{(\mathfrak{p})} \cong S(n)_{(g(\mathfrak{p}))} \quad (3.13)$$

for any $g \in \mathrm{SL}(2)$. In other words, we get a natural $\mathrm{SL}(2)$ -equivariant structure on $\mathcal{O}_{\mathbb{P}^1}(n)$ for any n ,

$$\Delta^n : q_{\mathrm{SL}(2)}^* \mathcal{O}_{\mathbb{P}^1}(n) \cong \delta^* \mathcal{O}_{\mathbb{P}^1}(n)$$

locally given by (3.13) where $\delta : \mathrm{SL}(2) \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is the action and $q_{\mathrm{SL}(2)} : \mathrm{SL}(2) \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is the second projection.

Let $\mathrm{SL}(2)$ act trivially on X , then $\mathrm{SL}(2)$ acts on $X \times \mathbb{P}^1$ by

$$g \circ (x, P) = (x, g \circ P)$$

for any $(x, P) \in X \times \mathbb{P}^1$. Under this action, p^*E_2 and p^*E_1 are $\mathrm{SL}(2)$ -invariant vector bundles and the morphism

$$p^*\varphi : p^*E_2 \rightarrow p^*E_1$$

is $\mathrm{SL}(2)$ -equivariant.

On \mathbb{P}^1 we have an exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^1}(-1) \rightarrow V \otimes \mathcal{O}_{\mathbb{P}^1} \xrightarrow{e_0} \mathcal{O}_{\mathbb{P}^1}(1) \rightarrow 0 \quad (3.14)$$

where e_0 is the evaluation map defined by

$$(e_0)_P(s \otimes \rho) = s_P \cdot \rho.$$

Here s_P is the image of global section $s \in V$ in the stalk $\mathcal{O}_{\mathbb{P}^1}(1)_P, P \in \mathbb{P}^1$ and $\rho \in \mathcal{O}_{\mathbb{P}^1, P}$. Moreover, for any $g \in \mathrm{SL}(2)$, $\Delta_{(g, P)}^1(s_P)$ is the image of the global section $g(s)$ in the stalk $\mathcal{O}_{\mathbb{P}^1}(1)_{g \circ P}$. So we have the following commutative diagram

$$\begin{array}{ccc} V \otimes \mathcal{O}_{\mathbb{P}^1, P} & \longrightarrow & \mathcal{O}_{\mathbb{P}^1}(1)_P \\ \downarrow g \otimes \Delta_{(g, P)} & & \downarrow \Delta_{(g, P)}^1 \\ V \otimes \mathcal{O}_{\mathbb{P}^1, g \circ P} & \longrightarrow & \mathcal{O}_{\mathbb{P}^1}(1)_{g \circ P} \end{array}$$

for any $g \in \mathrm{SL}(2)$ and $P \in \mathbb{P}^1$. It implies that e_0 is a $\mathrm{SL}(2)$ -equivariant morphism and hence $\mathcal{O}_{\mathbb{P}^1}(-1)$ is $\mathrm{SL}(2)$ -invariant. As $e = e_0 \otimes \mathrm{Id}_{\mathcal{O}_{\mathbb{P}^1}(-1)}$, the Euler sequence (3.9) is $\mathrm{SL}(2)$ -equivariant. It follows that the morphism

$$\psi = \mathrm{Id}_{p^*E_1} \otimes q^*e : p^*E_1 \otimes V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) \rightarrow p^*E_1$$

is $\mathrm{SL}(2)$ -equivariant. Being the fullback of two $\mathrm{SL}(2)$ -equivariant morphisms, $p^*\varphi$ and ψ , E is also $\mathrm{SL}(2)$ -invariant.

Proposition 3.3.2. *Let E be a vector bundle on X and $F \subseteq p^*E$ be a subbundle of p^*E on $X \times \mathbb{P}^1$ which is $\mathrm{SL}(2)$ -invariant. Then $F \cong p^*E'$ for some subbundle $E' \subseteq E$.*

Proof. To prove, we will show that $F|_{F_p} \cong \mathcal{O}_{\mathbb{P}^1}^{\oplus r}$ for any p -fiber F_p . Then Lemma 2.2.2 implies that $F \cong p^*E'$ for some $E' \subseteq E$. As a consequence of Proposition 1.1.9, there exists a filtration

$$0 = E_0 \subseteq E_1 \subseteq \dots \subseteq E_r = E$$

of subbundles such that E_i/E_{i-1} is a line bundle for each $i = 1, \dots, r$. Pulling back this sequence by p gives us a filtration for p^*E ,

$$0 = p^*E_0 \subseteq p^*E_1 \subseteq \dots \subseteq p^*E_r = p^*E,$$

with the same property. We intersect this sequence with F to get a filtration for F ,

$$0 = F_0 \subseteq F_1 \subseteq \dots \subseteq F_{r'} = F,$$

where $F_i = p^*E_i \cap F$, $r' = \mathrm{rank}(F)$ such that F_i/F_{i-1} is a line bundle. For each $i = 1, \dots, r'$, we consider the following $\mathrm{SL}(2)$ -equivariant diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & F_i & \longrightarrow & F_{i+1} & \longrightarrow & F_{i+1}/F_i \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & p^*E_i & \longrightarrow & p^*E_{i+1} & \longrightarrow & p^*E_{i+1}/p^*E_i \longrightarrow 0 \end{array}$$

Restricting this diagram to $F_p \cong \mathbb{P}^1$, we get

$$\begin{array}{ccccccc} 0 & \longrightarrow & F_i|_{F_p} & \longrightarrow & F_{i+1}|_{F_p} & \longrightarrow & \mathcal{O}_{\mathbb{P}^1}(a_i) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & p^*E_i|_{F_p} & \longrightarrow & p^*E_{i+1}|_{F_p} & \longrightarrow & \mathcal{O}_{\mathbb{P}^1} \longrightarrow 0 \end{array}$$

which is still a $\mathrm{SL}(2)$ -equivariant diagram. Consider the $\mathrm{SL}(2)$ -equivariant short exact sequences

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^1}(a_i) \longrightarrow \mathcal{O}_{\mathbb{P}^1} \longrightarrow T_i \longrightarrow 0 \quad (3.15)$$

where T_i is torsion sheaf for $i = 1, \dots, r'$. Since $\mathcal{O}_{\mathbb{P}^1}$ and $\mathcal{O}_{\mathbb{P}^1}(a_i)$ are $\mathrm{SL}(2)$ -invariant, T is also $\mathrm{SL}(2)$ -invariant. Moreover, $\mathrm{SL}(2)$ acts transitively on \mathbb{P}^1 . It follows that $\mathrm{Supp} T_i$ is whole \mathbb{P}^1 if it is non-empty. This contradicts the fact that T_i is torsion. Hence $T_i = 0$ and $\mathcal{O}_{\mathbb{P}^1}(a_i) \cong \mathcal{O}_{\mathbb{P}^1}$. But $F|_{F_p} \cong \bigoplus_{i=1}^{r'} \mathcal{O}_{\mathbb{P}^1}(a_i)$, therefore

$$F|_{F_p} \cong \bigoplus_{i=1}^{r'} \mathcal{O}_{\mathbb{P}^1}.$$

□

We get immediately the following corollary.

Corollary 3.3.3. *Let E_T be the vector bundle on $X \times \mathbb{P}^1$ defined by triple $T = (E_2 \xrightarrow{\varphi} E_1)$ on X and $E' \subset E_T$ be a $\mathrm{SL}(2)$ -invariant subbundle. Then E' is defined by a subtriple T' of T*

Proof. Let K and I be the kernel and image of morphism $E' \rightarrow E_T \rightarrow p^*E_2$. Then K and I are $\mathrm{SL}(2)$ -invariant bundles making the following diagram commutative

$$\begin{array}{ccccccc} 0 & \longrightarrow & K & \longrightarrow & E' & \longrightarrow & I \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2) & \longrightarrow & E_T & \longrightarrow & p^*E_2 \longrightarrow 0. \end{array} \quad (3.16)$$

By Proposition 3.3.2, $I = p^*E'_2$ and $K = p^*E'_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2)$ for some $E'_1 \subset E_1$ and $E'_2 \subset E_2$. Taking the pushforward of the diagram (3.16) by p , we get a commutative square

$$\begin{array}{ccc} E'_2 & \longrightarrow & E'_1 \otimes R^1p_*(q^*\mathcal{O}_{\mathbb{P}^1}(-2)) \\ \downarrow & & \downarrow \\ E_2 & \longrightarrow & E_1 \otimes R^1p_*(q^*\mathcal{O}_{\mathbb{P}^1}(-2)). \end{array}$$

By Serre's duality, we can take two isomorphisms

$$E'_1 \otimes R^1p_*(q^*\mathcal{O}_{\mathbb{P}^1}(-2)) \cong E'_1,$$

$$E_1 \otimes R^1p_*(q^*\mathcal{O}_{\mathbb{P}^1}(-2)) \cong E_1$$

such that the diagram

$$\begin{array}{ccc} E'_2 & \xrightarrow{\varphi'} & E'_1 \\ \downarrow & & \downarrow \\ E_2 & \xrightarrow{\varphi} & E_1 \end{array}$$

commutes, i.e. $T' = (E'_2 \xrightarrow{\varphi'} E'_1)$ is a subtriple of T . □

For a given triple $T = (E_2, E_1, \varphi)$ of type (r_1, r_2, d_1, d_2) , we set

$$\alpha_m = \frac{\mu(E_1) - \mu(E_2)}{2} \text{ and } \alpha_M = \left(1 + \frac{r_1 + r_2}{|r_1 - r_2|}\right) \frac{\mu(E_1) - \mu(E_2)}{2} \quad (3.17)$$

if $r_1 \neq r_2$ and $\alpha_M = +\infty$ if $r_1 = r_2$. We have seen from Propositions 3.2.2 and 3.2.3 that T can not be α -semistable unless

$$\alpha_m \leq \alpha \leq \alpha_M \text{ and } \alpha \geq 0.$$

Let $\alpha = \frac{a}{b} \in \mathbb{Q}$ such that $a, b > 0$, $\gcd(a, b) = 1$ and $\alpha_m \leq \alpha \leq \alpha_M$. Let E_T be the vector bundle on $X \times \mathbb{P}^1$ defined by T as in (3.11) and (3.12). It is obvious that

$$E_T \in \text{Ext}^1(p^*E_2, p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2)).$$

On $X \times \mathbb{P}^1$, $H(\alpha) = aF_p + bF_q$ is an ample divisor. We have

$$\deg_{H(\alpha)}(E_T) = b(d_1 + d_2) - 2ar_1$$

and then the $H(\alpha)$ -slope of E_T

$$\mu_{H(\alpha)}(E_T) = \frac{b(d_1 + d_2) - 2ar_1}{r_1 + r_2} = b\mu_\alpha(T).$$

Theorem 3.3.4. *The triple $T = (E_2, E_1, \varphi)$ is α -(semi)stable if and only if the extension E_T defined by T is $H(\alpha)$ -(semi)stable.*

Proof. Assume that the bundle E_T defined by T is $H(\alpha)$ -(semi)stable. Let $T' = (E'_2 \xrightarrow{\varphi'} E'_1)$ be a nontrivial subtriple of T . It follows from Lemma 3.3.1 that the bundle $E'_{T'}$ defined by T' is a subbundle of E_T . Hence

$$\mu_\alpha(T') = \frac{\mu_{H(\alpha)}(E'_{T'})}{b} (\leq) \frac{\mu_{H(\alpha)}(E_T)}{b} = \mu_\alpha(T),$$

i.e. T is α -(semi)stable.

Assume that T is α -(semi)stable and E_T is not $H(\alpha)$ -(semi)stable. Then there exists a maximal destabilizing subsheaf $E' \subseteq E_T$. As E_T is $\text{SL}(2)$ -invariant, E' is also $\text{SL}(2)$ -invariant because $g^*E' \subseteq E_T$ for every $g \in \text{SL}(2)$ and the action of g does not change the rank E' and the $\deg_{H(\alpha)}(E')$ as $\text{SL}(2)$ is connected. By Corollary 3.3.2, $E' = E'_T$ for some subtriple $T' = (E'_2 \xrightarrow{\varphi'} E'_1)$ of T . We have

$$\mu_\alpha(T') = \frac{\mu_{H(\alpha)}(E'_T)}{b} (\geq) \frac{\mu_{H(\alpha)}(E_T)}{b} = \mu_\alpha(T).$$

This contradicts the α -(semi)stability of T . □

3.4 Moduli space of triples

Let S be a scheme of finite type over k and $p : S \times X \rightarrow S$ be the first projection. A family of triples on X parametrized by S (or a S -family of triples on X) is a triple $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ on $S \times X$ where \mathcal{E}_i are families of vector bundles on X parametrized by S . Let $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ and $\mathcal{T}' = (\mathcal{E}'_2 \xrightarrow{\varphi'_S} \mathcal{E}'_1)$ be S -families of triples. We say that they are equivalent if there exists a line bundle L on S and isomorphisms $\psi_i : \mathcal{E}_i \rightarrow \mathcal{E}'_i \otimes p^*L$ such that

$$\varphi_S = \psi_1^{-1} \circ (\varphi'_S \otimes p^* \text{Id}_L) \circ \psi_2.$$

For a given $\alpha \in \mathbb{Q}_{>0}$ we say that $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ is a S -family of α -(semi)stable triples of type $(\underline{r}, \underline{d}) = (r_1, r_2, d_1, d_2)$ if $\mathcal{T}_s = (\mathcal{E}_{2s} \xrightarrow{\varphi_s} \mathcal{E}_{1s})$ is α -(semi)stable and \mathcal{E}_{is} have rank r_i and degree $d_i, i = 1, 2; s \in S$. Let Sch/k and $Sets$ be the category of schemes of finite type over k and the category of sets, respectively. We consider the following contravariant functor:

$$\begin{aligned} \underline{\mathcal{M}}_\alpha^{(s)s}(\underline{r}, \underline{d}) : Sch/k &\longrightarrow Sets \\ S &\longmapsto \{\text{Equivalence classes of } S - \text{families of } \alpha - (\text{semi})\text{stable} \\ &\quad \text{triples of type } (\underline{r}, \underline{d})\}. \end{aligned}$$

There are several approaches to construct the moduli space of α -(semi)stable triples on curves, i.e. the scheme corepresents $\underline{\mathcal{M}}_\alpha^{(s)s}(\underline{r}, \underline{d})$. As we have seen, a triple $T = (E_2 \xrightarrow{\varphi} E_1)$ is α -(semi)stable if and only if the vector bundle E_T on the ruled surface $X \times \mathbb{P}^1$ defined by T is $H(\alpha)$ -(semi)stable. It implies that we can embed the functor $\underline{\mathcal{M}}_\alpha^{(s)s}(\underline{r}, \underline{d})$ into the moduli functor of $H(\alpha)$ -(semi)stable vector bundles on $X \times \mathbb{P}^1$ which is corepresented by a projective scheme of finite type over k (cf. [13], [18]). By this way, Brallow and García-Prada (cf. [2]) constructed the moduli space of α -stable triples on compact Riemann surfaces.

From another point of view, any triple $T = (E_2 \xrightarrow{\varphi} E_1)$ on X can be seen as a representation of the quiver Q ,

$$\bullet \longrightarrow \bullet,$$

consisting of 2 vertices and one arrow, in the category of vector bundles on X . As the semistability of representations is given, we also have the concepts of the Harder-Narasimhan filtration, Jordan-Hölder filtration, the associated graded object for a representation and then the S -equivalence as usual. The moduli space of α -semistable triples is then just a special case of the moduli space of twisted representation of quivers given by A.Schmitt (cf. [17, Thm. 3.7.2]) as the base field k is the field of complex numbers or by Álavrez-Cónsul (cf. [1]) for characteristic p . From there, we obtain the moduli space of α -semistable triples on smooth projective curve X .

Theorem 3.4.1. *There exist a projective scheme $\mathcal{M}_\alpha^{ss}(\underline{r}, \underline{d})$ which corepresents the functor $\underline{\mathcal{M}}_\alpha^{ss}(\underline{r}, \underline{d})$. The closed points of $\mathcal{M}_\alpha^{ss}(\underline{r}, \underline{d})$ are in bijection to the set of S -equivalence classes of α -semistable triples of type (r_1, r_2, d_1, d_2) . There is an open subscheme $\mathcal{M}_\alpha^s(\underline{r}, \underline{d})$ which corepresents the subfunctor $\underline{\mathcal{M}}_\alpha^s(\underline{r}, \underline{d})$ and whose closed points correspond to the isomorphism classes of α -stable triples.*

Proof. See [17, Thm. 3.7.2] or [1, Thm. 8] □

Chapter 4

Generalized theta line bundles

4.1 Theta divisors for vector bundles on curves

4.1.1 Determinant line bundles

Let X be a smooth projective variety of dimension n . We define the Grothendieck's groups $K(X)$ and $K^0(X)$ to be the quotients of free abelian group generated by coherent and locally free sheaves, respectively, by the subgroup generated by all expressions $E - E' - E''$, wherever there is an exact sequence $0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0$. Moreover, the tensor product turns $K^0(X)$ into a commutative ring with $1 = [\mathcal{O}_X]$ and gives $K(X)$ a module structure over $K^0(X)$.

Let S be a connected scheme of finite type over k . Then the projection $p : S \times X \rightarrow S$ is a smooth morphism of relative dimension n . Let \mathcal{E} be a flat family of coherent sheaves on X parametrized by S . We have a homomorphism $p_! : K^0(S \times X) \rightarrow K^0(S)$, given by

$$p_!([\mathcal{E}]) = \sum_{i=0}^n (-1)^i [R^i p_* \mathcal{E}]$$

(cf. [12, Cor. 2.1.11]). Let $q : S \times X \rightarrow X$ be the second projection. Then we have a well defined morphism

$$\begin{aligned} \lambda_{\mathcal{E}} : K(X) &\rightarrow \text{Pic}(S) \\ [F] &\mapsto \det(p_!(\mathcal{E} \otimes q^*[F])) \end{aligned} \tag{4.1}$$

which is the composition of morphisms

$$K(X) \xrightarrow{q^*} K^0(S \times X) \xrightarrow{[\cdot \mathcal{E}]} K^0(S \times X) \xrightarrow{p_!} K^0(S) \xrightarrow{\det} \text{Pic}(S).$$

For each class $[F] \in K(X)$, the line bundle $\lambda_{\mathcal{E}}([F])$ is called a determinant line bundle associated to \mathcal{E} at the class $[F]$. The following properties of $\lambda_{\mathcal{E}}$ can be seen easily by using

the projection formula for direct images of sheaves and the basic properties of the determinant of locally free sheaves.

Lemma 4.1.1 (cf. [12, Lm. 8.1.2]). *Let \mathcal{E} be a S -flat family of coherent sheaves and $[F]$ be any class in $K(X)$.*

(i) *(Additive property) If*

$$0 \longrightarrow \mathcal{E}' \longrightarrow \mathcal{E} \longrightarrow \mathcal{E}'' \longrightarrow 0$$

is a short exact sequence of S -flat families of coherent sheaves then

$$\lambda_{\mathcal{E}}([F]) = \lambda_{\mathcal{E}'}([F]) \otimes \lambda_{\mathcal{E}''}([F]).$$

Similarly, if we have a short exact sequence of coherent sheaves on X

$$0 \longrightarrow F' \longrightarrow F \longrightarrow F'' \longrightarrow 0,$$

then $\lambda_{\mathcal{E}}([F]) \cong \lambda_{\mathcal{E}}([F']) \otimes \lambda_{\mathcal{E}}([F''])$.

(ii) *(Base-change property) If $f : S' \longrightarrow S$ is any morphism, then*

$$f^*(\lambda_{\mathcal{E}}([F])) = \lambda_{f^*\mathcal{E}}([F])$$

where $f_X := f \times \text{Id}_X : S' \times X \longrightarrow S \times X$.

(iii) *(Projection property) If L is a line bundle on S then*

$$\lambda_{\mathcal{E} \otimes p^*L}([F]) = \lambda_{\mathcal{E}}([F]) \otimes L^{\chi(\mathcal{E}_s \otimes F)}$$

for any $s \in S$.

4.1.2 Generalized theta functions and generalized theta divisors

Let X be a smooth projective curve and r, d be integers, $r \geq 2$. Let $U_X(r, d)$ be the moduli space of S -equivalence classes of vector bundles on X of rank r and degree d . Then $U_X(r, d)$ is a normal projective variety whose smooth locus is the fine moduli space of stable vector bundles $U_X^s(r, d)$, unless $g = 2, r = 2$ and d is even, when $U_X(r, d)$ is smooth. Drezet and Narasimhan (cf. [5]) even showed that, they are locally factorial schemes. Let F be a nonzero vector bundle on X . We call F a complementary vector bundle of $U_X(r, d)$ if $\chi(E \otimes F) = 0$ for all $E \in U_X(r, d)$. Let $h = \gcd(r, d)$ and $r_0 = \frac{r}{h}, d_0 = \frac{d}{h}$. By Riemann-Roch formula, it is easy to see that F has rank pr_0 and degree $p(r_0(g_X - 1) - d_0)$ for some $p \in \mathbb{Z}_{>0}$. In particular, to a generic vector bundle F_0 of rank r_0 and degree $r_0(g_X - 1) - d_0$ we can associate an effective Cartier divisor Θ_{F_0} on $U_X(r, d)$ (cf. [5, Section 0.2]) supported on the set

$$\Theta_{F_0} = \{E \in U_X(r, d) \mid H^0(E \otimes F) \neq 0\},$$

called a basic theta divisor.

Let S be a connected scheme and \mathcal{E} be a family of vector bundles of rank r and degree d parametrized by S . We will show by the following construction that for each vector bundle F on X such that $\chi(\mathcal{E}_s \otimes F) = 0$ for some $s \in S$ (hence for all $s \in S$), the line bundle $\lambda_{\mathcal{E}}([F])^{-1}$ has a geometric section $\zeta_{\mathcal{E}}(F)$ whose zero divisor is given by

$$\{s \in S \mid H^0(X, \mathcal{E}_s \otimes F) \neq 0\}. \quad (4.2)$$

Indeed, there exists $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$,

$$H^0(X, \mathcal{E}_s \otimes \mathcal{O}_X(-n)) = 0$$

for all $s \in S$. Let F be any vector bundle of rank r' on X . Proposition 1.1.9 implies that for n large enough, there exists a short exact sequence of vector bundles on X

$$0 \longrightarrow \det(F)^{-1}(-(1+r')n) \xrightarrow{\varphi_F(n)} \mathcal{O}_X^{r'+1}(-n) \longrightarrow F \longrightarrow 0. \quad (4.3)$$

We consider such large n that it is also bigger than n_0 and denote by $n \gg n_0$. Pulling back this exact sequence by q and then tensoring with \mathcal{E} , we get an exact sequence on $S \times X$

$$0 \longrightarrow \mathcal{E} \otimes q^* \det(F)^{-1}(-(1+r')n) \xrightarrow{\text{Id}_{\mathcal{E}} \otimes q^* \varphi_F(n)} \mathcal{E} \otimes q^* \mathcal{O}_X^{r'+1}(-n) \longrightarrow \mathcal{E} \otimes q^* F \longrightarrow 0.$$

Pushing forward to S by p , we have

$$\begin{aligned} 0 \longrightarrow p_*(\mathcal{E} \otimes q^* \det(F)^{-1}(-(r'+1)n)) &\longrightarrow p_*(\mathcal{E} \otimes q^* \mathcal{O}_X^{r'+1}(-n)) \longrightarrow p_*(\mathcal{E} \otimes q^* F) \\ &\longrightarrow R^1 p_*(\mathcal{E} \otimes q^* \det(F)^{-1}(-(r'+1)n)) \xrightarrow{R^1(\varphi_F(n))} R^1 p_*(\mathcal{E} \otimes q^* \mathcal{O}_X^{r'+1}(-n)) \\ &\longrightarrow R^1 p_*(\mathcal{E} \otimes q^* F) \longrightarrow 0 \end{aligned} \quad (4.4)$$

where $R^1(\varphi_F(n)) = R^1 p_*(\text{Id}_{\mathcal{E}} \otimes q^* \varphi_F(n))$. For any $s \in S$, we have

$$p_*(\mathcal{E} \otimes q^* \mathcal{O}_X^{r'+1}(-n)) \otimes k(s) \cong H^0(X, \mathcal{E}_s^{r'+1}(-n)) = 0.$$

Hence the exact sequence (4.4) becomes

$$\begin{aligned} 0 \longrightarrow p_*(\mathcal{E} \otimes q^* F) &\longrightarrow R^1 p_*(\mathcal{E} \otimes q^* \det(F)^{-1}(-(r'+1)n)) \xrightarrow{R^1(\varphi_F(n))} R^1 p_*(\mathcal{E} \otimes q^* \mathcal{O}_X^{r'+1}(-n)) \\ &\longrightarrow R^1 p_*(\mathcal{E} \otimes q^* F) \longrightarrow 0. \end{aligned} \quad (4.5)$$

Then we have

$$\lambda_{\mathcal{E}}([F])^{-1} \cong \det(R^1 p_*(\mathcal{E} \otimes q^* \det(F)^{-1}(-(r'+1)n)))^{-1} \otimes \det(R^1 p_*(\mathcal{E} \otimes q^* \mathcal{O}_X^{r'+1}(-n))). \quad (4.6)$$

Assume that for some $m \gg n_0$, we have another resolution of F of form

$$0 \longrightarrow \det(F)^{-1}(-(1+r')m) \xrightarrow{\varphi_F(m)} \mathcal{O}_X^{r'+1}(-m) \longrightarrow F \longrightarrow 0. \quad (4.7)$$

Then we have the same long exact sequence as (4.5) for m . It implies that

$$\begin{aligned} & \det(R^1 p_*(\mathcal{E} \otimes q^* \det(F)^{-1}(-(r' + 1)n)))^{-1} \otimes \det(R^1 p_*(\mathcal{E} \otimes q^* \mathcal{O}_X^{r'+1}(-n))) \\ & \cong \det(R^1 p_*(\mathcal{E} \otimes q^* \det(F)^{-1}(-(r' + 1)m)))^{-1} \otimes \det(R^1 p_*(\mathcal{E} \otimes q^* \mathcal{O}_X^{r'+1}(-m))), \end{aligned} \quad (4.8)$$

equivalently, $\lambda_{\mathcal{E}}([F])^{-1}$ is independent of the choice of n provided $n \gg n_0$. It is then clear that

$$\lambda_{\mathcal{E}}([F]) \cong \lambda_{\mathcal{E}}([F']) \quad (4.9)$$

for any F, F' such that $\text{rank}(F) = \text{rank}(F')$ and $\det(F) = \det(F')$.

Assume that $\chi(\mathcal{E}_s \otimes F) = 0$ for some $s \in S$ (hence for all $s \in S$). It follows from Riemann - Roch formula that

$$\text{rank}(p_*(\mathcal{E} \otimes q^* F)) = \text{rank}(R^1 p_*(\mathcal{E} \otimes q^* F)).$$

From the exactness of (4.5) we have

$$\text{rank}(R^1 p_*(\mathcal{E} \otimes q^* \det(F)^{-1}(-(r' + 1)n))) = \text{rank}(R^1 p_*(\mathcal{E} \otimes q^* \mathcal{O}_X^{r'+1}(-n))).$$

It implies that

$$R^1(\varphi_F(n)) : R^1 p_*(\mathcal{E} \otimes q^* \det(F)^{-1}(-(r' + 1)n)) \longrightarrow R^1 p_*(\mathcal{E} \otimes q^* \mathcal{O}_X^{r'+1}(-n))$$

is a morphism of vector bundles of the same rank. Therefore the line bundle

$$\lambda_{\mathcal{E}}([F])^{-1} \cong \det(R^1 p_*(\mathcal{E} \otimes q^* \det(F)^{-1}(-(r' + 1)n)))^{-1} \otimes \det(R^1 p_*(\mathcal{E} \otimes q^* \mathcal{O}_X^{r'+1}(-n)))$$

has a geometric section $\zeta_{\mathcal{E}}(F) = \det(R^1(\varphi_F(n)))$. Since $\chi(\mathcal{E}_s \otimes F) = 0$ for all $s \in S$, by the construction, we see that the zero divisor of $\zeta_{\mathcal{E}}(F)$ parametrizes all points $s \in S$ such that $H^0(S, \mathcal{E}_s \otimes F) \neq 0$ (cf. [11, Prop. 4.1]).

Assume further that \mathcal{E} is a family of semistable vector bundles of rank r and degree d on X parametrized by S . Then there exists a morphism

$$f_{\mathcal{E}} : S \longrightarrow U_X(r, d), s \mapsto \mathcal{E}_s$$

induced by the family \mathcal{E} . Let $[F] \in K(X)$ be any class where F is a complementary vector bundle of $U_X(r, d)$. Then $\chi(\mathcal{E}_s \otimes F) = 0$ for all $s \in S$. We obtain on S a line bundle $\lambda_{\mathcal{E}}([F])^{-1}$ and a generalized theta function $\zeta_{\mathcal{E}}(F)$.

Theorem 4.1.2 (cf. [5, Thm. D, part (a)], [15, Cor. 1.6, Cor. 1.7] or [12, Thm. 8.1.5]). *Let $[F]$ be a class in $K(X)$ such that F is a complementary vector bundle of $U_X(r, d)$. There exists a unique ample line bundle $\mathbb{L}([F])$ on $U_X(r, d)$ such that for any family \mathcal{E} of semistable vector bundles of rank r and degree d on X parametrized by S , we have*

$$\lambda_{\mathcal{E}}([F])^{-1} \cong f_{\mathcal{E}}^* \mathbb{L}([F]). \quad (4.10)$$

Moreover, the geometric section $\zeta_{\mathcal{E}}(F)$ descends by $f_{\mathcal{E}}$ to a section of $\mathbb{L}([F])$ on $U_X(r, d)$ whose zero divisor is supported on the set

$$\Theta_F = \{E \in U_X(r, d) \mid H^0(X, E \otimes F) \neq 0\}.$$

We denote this section by ζ_F and call it a generalized theta function. As $\mathbb{L}([F])$ is independent of the choice of representations of class $[F] \in K(X)$, we denote it again by θ_F and call it a generalized theta line bundle. If ζ_F is a nonzero section, i.e. there exists $E \in U_X(r, d)$ such that $H^0(E \otimes F) = 0$, then Θ_F is a divisor, called a generalized theta divisor, and $\theta_F = \mathcal{O}_{U_X(r, d)}(\Theta_F)$. In particular, if we take $F = F_0$ then $\lambda_{\mathcal{E}}([F_0])^{-1}$ descends to basic theta line bundle $\theta_{F_0} = \mathcal{O}_{U_X(r, d)}(\Theta_{F_0})$. Let F'_0 be another vector bundle such that $\Theta_{F'_0}$ is a basic theta divisor. We have a morphism of Picard groups $\det^* : \text{Pic}(J^d(X)) \rightarrow \text{Pic}(U_X(r, d))$ induced from the canonical morphism $\det : U_X(r, d) \rightarrow J^{(d)}(X)$. As $\text{Pic}^0(X) \subset \text{Pic}(J^d(X))$, we have

$$\theta_{F'_0} \cong \theta_{F_0} \otimes \det^*(\det(F'_0) \otimes \det(F_0)^{-1}) \quad (4.11)$$

(cf. [5, Thm. D]).

4.1.3 Pluritheta linear series on $U_X(r, d)$

Proposition 1.1.6 gives us a sufficient condition under which a vector bundle is semistable: E is semistable if there exists a nonzero vector bundle F such that $E \otimes F$ is cohomologically trivial, i.e. $H^*(X, E \otimes F) = 0$. In fact, it was proven by Faltings in [8], Seshadri in [18, Thm. 6.2], etc, that this is also a necessary condition for semistability of vector bundles on curves. In [6, Thm. 2], Esteves even showed that we can find such vector bundle F with fixed determinant. We recall this result for $E \in U_X(r, d)$.

Theorem 4.1.3 ([6, Thm. 2]). *Let $E \in U_X(r, d)$ be a semistable vector bundle on X and $\{L_p \mid p > 0\}$ be a sequence of line bundles such that $\deg(L_p) = p(r_0(g_X - 1) - d_0)$ for every integer $p > 0$. Then there is a vector bundle F on X of rank pr_0 such that $\det(F) \cong L_p$ and $E \otimes F$ is cohomologically trivial for a certain $p > 0$.*

Let F_0 be a complementary vector bundle of $U_X(r, d)$ such that Θ_{F_0} is a basic theta divisor. For each integer p , let us set $L_p = (\det(F_0))^p$ and consider the complementary vector bundles F of rank pr_0 and determinant L_p . It follows from Theorem 4.1.2 and (4.11) that for any such F , the generalized theta line bundle θ_F is ample and isomorphic to $\theta_{F_0}^p$. In particular, if ζ_F is a nonzero section, then $\Theta_F \in |p\Theta_{F_0}|$. We call $|p\Theta_{F_0}|$ a pluritheta linear series. What we are interested in are the properties of this linear series as base point freeness, separated properties, very ampleness, etc. These properties should be related to a bound on the rank of F .

Theorem 4.1.4 ([7, Thm. B]). *Let Θ_{F_0} be a basic theta divisor on $U_X(r, d)$. For each $p \geq r^2 + r$, the linear series $|p\Theta_{F_0}|$ on $U_X(r, d)$ separates points and is very ample on the smooth locus $U_X^s(r, d)$.*

4.2 Determinant line bundles and invariant sections for triples

4.2.1 Triple product

For any triples $T = (E_2 \xrightarrow{\varphi} E_1)$ and $F = (F_2 \xrightarrow{\psi} F_1)$ on X , we define the triple product of T and F to be the triple

$$T \times F := (E_2 \otimes F_1 \oplus E_1 \otimes F_2 \xrightarrow{(\varphi \otimes \text{Id}_{F_1})\pi_1 - (\text{Id}_{E_1} \otimes \psi)\pi_2} E_1 \otimes F_1) \quad (4.12)$$

where $\pi_1 : E_2 \otimes F_1 \oplus E_1 \otimes F_2 \longrightarrow E_2 \otimes F_1$ and $\pi_2 : E_2 \otimes F_1 \oplus E_1 \otimes F_2 \longrightarrow E_1 \otimes F_2$ are the projections. It is easy to see that the triple product is exact, i.e. if

$$0 \longrightarrow T' \longrightarrow T \longrightarrow T'' \longrightarrow 0 \quad (4.13)$$

is an exact sequence of triples on X then for any triple F , the sequence

$$0 \longrightarrow T' \times F \longrightarrow T \times F \longrightarrow T'' \times F \longrightarrow 0$$

is exact.

Definition 4.2.1. *For a given triple $T = (E_2 \xrightarrow{\varphi} E_1)$ on X , we call $\chi(T)$ the Euler Characteristic of T , defined by*

$$\chi(T) = \chi(E_2) - \chi(E_1),$$

where $\chi(E_i)$ is the Euler Characteristic of $E_i, i = 1, 2$.

For an exact sequence of triples as (4.13), we have

$$\chi(T) = \chi(T') + \chi(T'').$$

4.2.2 α -orthogonal triples

Let $T = (E_2 \xrightarrow{\varphi} E_1)$ and $F = (F_2 \xrightarrow{\psi} F_1)$ be triples on X . Let K be the kernel of

$$\varphi \times \psi := (\varphi \otimes \text{Id}_{F_1})\pi_1 - (\text{Id}_{E_1} \otimes \psi)\pi_2$$

as in (4.12). We then have a short exact sequence

$$0 \longrightarrow K \longrightarrow E_2 \otimes F_1 \oplus E_1 \otimes F_2 \xrightarrow{\varphi \times \psi} E_1 \otimes F_1.$$

Assume that $T = (E_2 \xrightarrow{\varphi} E_1)$ is of type (r_1, r_2, d_1, d_2) . As in (3.17), let us set

$$\alpha_m = \frac{\mu(E_1) - \mu(E_2)}{2} \text{ and } \alpha_M = \left(1 + \frac{r_1 + r_2}{|r_1 - r_2|}\right) \frac{\mu(E_1) - \mu(E_2)}{2}$$

if $r_1 \neq r_2$ and $\alpha_M = +\infty$ if $r_1 = r_2$. From now on, by a rational number $\alpha = \frac{a}{b} \in \mathbb{Q}$, we mean a rational number $\alpha = \frac{a}{b} \in \mathbb{Q}$ such that $a, b > 0$, $(a, b) = 1$ and $\alpha_m \leq \alpha \leq \alpha_M$.

Definition 4.2.2. Let $T = (E_2 \xrightarrow{\varphi} E_1)$, $F = (F_2 \xrightarrow{\psi} F_1)$ be triples on X and $\alpha = \frac{a}{b} \in \mathbb{Q}$. Then F is called a α -orthogonal triple of T if it satisfies the following properties:

- (i) ψ is surjective,
- (ii) $\text{rank}(F_2) = 2\text{rank}(F_1)$, $\mu(F_2) = \mu(F_1) - \alpha$,
- (iii) $H^*(X, K) = 0$ where K is the kernel of $\varphi \times \psi$.

Similar to the semistability of vector bundles, we can characterize the α -semistability of triples by a perpendicularity condition.

Theorem 4.2.3. Let $T = (E_2 \xrightarrow{\varphi} E_1)$ be a triple of type (r_1, r_2, d_1, d_2) on X and $\alpha = \frac{a}{b} \in \mathbb{Q}$. Then T is α -semistable if and only if it has a α -orthogonal triple.

Proof. Assume that $F = (F_2 \xrightarrow{\psi} F_1)$ is a α -orthogonal triple of $T = (E_2 \xrightarrow{\varphi} E_1)$. If T is not α -semistable, there exist a subtriple $T' = (E'_2 \xrightarrow{\varphi'} E'_1)$ which destabilizes T , i.e. $\mu_\alpha(T') > \mu_\alpha(T)$. Consider the following commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & K' & \longrightarrow & E'_2 \otimes F_1 \oplus E'_1 \otimes F_2 & \longrightarrow & E'_1 \otimes F_1 & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & K & \longrightarrow & E_2 \otimes F_1 \oplus E_1 \otimes F_2 & \longrightarrow & E_1 \otimes F_1 & \longrightarrow & 0 \end{array}$$

where the rows are exact.

Assume that F is of type (R_1, R_2, D_1, D_2) . Then

$$R_2 = 2R_1, D_2 = 2D_1 - 2\alpha R_1$$

by condition (ii) in Definition 4.2.2. We have

$$\text{rank}(K) = (r_1 + r_2)R_1 \text{ and } \deg(K) = (r_1 + r_2)D_1 + R_1(d_1 + d_2) - 2\alpha r_1 R_1.$$

It follows that

$$\mu(K) = \mu_\alpha(T) + \mu(F_1). \tag{4.14}$$

Similarly, $\mu(K') = \mu_\alpha(T') + \mu(F_1)$. Since $H^*(X, K) = 0$ then $\chi(K) = 0$. Moreover, $K' \subseteq K$ implies that $\chi(K') \leq 0$. Therefore

$$\mu(K) = g_X - 1 \geq \mu(K') \implies \mu_\alpha(T) \geq \mu_\alpha(T').$$

This is a contradiction. Hence T is α -semistable.

Assume that T is α -semistable. We will construct a α -orthogonal triple $F = (F_2 \xrightarrow{\psi} F_1)$ of T . This is done by the following lemma.

Lemma 4.2.4. *Let $T = (E_2 \xrightarrow{\varphi} E_1)$ be a triple on X of type (r_1, r_2, d_1, d_2) and $\alpha = \frac{a}{b} \in \mathbb{Q}$. If T is α -semistable then it has a α -orthogonal triple.*

Proof. Let E_T be the extension on $X \times \mathbb{P}^1$ defined by T . It follows from Theorem 3.3.4 that E_T is $H(\alpha)$ -semistable of rank $r = r_1 + r_2$ and degree $b(d_1 + d_2) - 2ar_1$. By computing the Chern character up to numerical equivalence, we have

$$\text{ch}(E_T) \equiv r_1 + r_1 + (d_1 + d_2)F_p - 2r_1F_q - 2d_1[pt].$$

Therefore

$$\Delta(E_T) = 2 \text{ch}_0(E_T) \text{ch}_2(E_T) - \text{ch}_1^2(E_T) = 4(r_1d_2 - r_2d_1). \quad (4.15)$$

Let us fix $C \in |m_0H(\alpha)|$ a smooth projective curve where

$$m_0 = \lceil \frac{1-r}{r} \Delta(E_T) + 1 \rceil = \lceil \frac{4(1-(r_1+r_2))(d_2r_1-d_1r_2)}{r_1+r_2} + 1 \rceil. \quad (4.16)$$

Here we denote by $\lceil x \rceil$ the smallest integral number which is not less than x for any $x \in \mathbb{Q}$. Applying Corollary 2.3.4 for E_T , we see that $E_T|_C$ is again a semistable vector bundle of rank r and degree

$$d = m_0 \deg_{H(\alpha)}(E_T) = bm_0\mu_\alpha(T).$$

By Theorem 4.1.3, there exists a vector bundle F_C on C such that $H^*(C, E_T|_C \otimes F_C) = 0$. Let f be the composition $C \hookrightarrow X \times \mathbb{P}^1 \rightarrow X$ which is a finite morphism. Consider the following diagram,

$$\begin{array}{ccccccc} 0 & \longrightarrow & p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2) & \longrightarrow & E_T & \longrightarrow & p^*E_2 \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2) & \longrightarrow & p^*E_1 \otimes V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1) & \longrightarrow & p^*E_1 \longrightarrow 0 \end{array}$$

By restricting the above diagram to C and then twisting with F_C , the square on the left hand side becomes

$$\begin{array}{ccc} E_T|_C \otimes F_C & \longrightarrow & p^*E_2|_C \otimes F_C \\ \downarrow & & \downarrow \\ [p^*E_1 \otimes V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)]|_C \otimes F_C & \longrightarrow & p^*E_1|_C \otimes F_C \end{array}$$

We take the pushforward of this diagram to X by f and set

$$K = f_*(E_T|_C \otimes F_C), F_1 = f_*F_C, F_2 = f_*((V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1))|_C \otimes F_C).$$

As f is finite,

$$R^1 f_*([p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2)]_C \otimes F_C) = 0.$$

So we get a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & f_*([p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2)]_C \otimes F_C) & \longrightarrow & K & \longrightarrow & E_2 \otimes F_1 \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \varphi \otimes \text{Id}_{F_1} \\ 0 & \longrightarrow & f_*([p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2)]_C \otimes F_C) & \longrightarrow & E_1 \otimes F_2 & \xrightarrow{\text{Id}_{F_1} \otimes \psi} & E_1 \otimes F_1 \longrightarrow 0, \end{array}$$

where $\psi := f_*(q^*e|_C \otimes \text{Id}_{F_C}) : F_2 \longrightarrow F_1$ is surjective. It implies that K is the pullback of two morphisms

$$\varphi \otimes \text{Id}_{F_1} : E_2 \otimes F_1 \longrightarrow E_1 \otimes F_1 \text{ and } \text{Id}_{E_1} \otimes \psi : E_1 \otimes F_2 \longrightarrow E_1 \otimes F_1.$$

Equivalently, K is the kernel of $\varphi \times \psi$, the morphism of triple product

$$(E_2 \xrightarrow{\varphi} E_1) \times (F_2 \xrightarrow{\psi} F_1).$$

We then have a short exact sequence of vector bundles,

$$0 \longrightarrow K \longrightarrow E_2 \otimes F_1 \oplus E_1 \otimes F_2 \xrightarrow{\varphi \times \psi} E_1 \otimes F_1 \longrightarrow 0.$$

Since f is finite,

$$H^*(X, K) = H^*(X, f_*(E_T|_C \otimes F_C)) = H^*(C, E_T|_C \otimes F_C) = 0.$$

So $F = (F_2 \xrightarrow{\psi} F_1)$ will be the triple which we are looking for if the numerical invariants of F_1 and F_2 satisfy condition (ii) in Definition 4.2.2.

Now we compute the rank and degree of $F_1 = f_*(F_C)$. Let us set

$$h = \gcd(r, d), r_0 = \frac{r}{h}, d_0 = \frac{d}{h}.$$

As we have seen,

$$\text{rank } F_C = pr_0, \deg(F_C) = p(r_0(g_C - 1) - d_0) \quad (4.17)$$

for some $p \in \mathbb{Z}_{>0}$. Let G be a vector bundle on C of rank R and degree D . Since $C \in |m_0 H(\alpha)|$ where $H(\alpha) = aF_p + bF_q$, $f : C \longrightarrow X$ is finite of degree $m_0 b$. For any $x \in X$,

$$\dim(f_*G \otimes k(x)) = m_0 b R.$$

Hence f_*G is a vector bundle of rank m_0bR . As f is finite, we have

$$\dim H^*(C, G) = \dim H^*(X, f_*G).$$

So Riemann-Roch formula implies that $\chi(f_*G) = \chi(G)$. But

$$\chi(f_*G) = \chi(G) \iff \deg(f_*G) + \text{rank}(f_*G)(1 - g_X) = D + R(1 - g_C).$$

it implies that $\deg(f_*G) = RB + D$ where $B = m_0b(g_X - 1) + 1 - g_C$. Now we have,

$$\begin{aligned} \text{rank}(F_1) &= m_0bpr_0, \\ \text{rank}(F_2) &= 2m_0bpr_0 = 2\text{rank}(F_1), \\ \deg(F_1) &= pr_0(m_0b(g_X - 1) + 1 - g_C) + p(r_0(g_C - 1) - d_0) \\ &= m_0bpr_0(g_X - 1) - pd_0, \\ \deg(F_2) &= 2pr_0(m_0b(g_X - 1) + 1 - g_C) + \deg(V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)|_C \otimes F_C) \\ &= 2pr_0(m_0b(g_X - 1) + 1 - g_C) - 2kr_0m_0a + 2p(r_0(g_C - 1) - d_0) \\ &= 2(m_0bpr_0(g_X - 1) - pd_0) - 2pr_0m_0a \\ &= 2\deg(F_1) - 2\alpha\text{rank}(F_1). \end{aligned}$$

It follows that

$$\text{rank}(F_2) = 2\text{rank}(F_1), \mu(F_2) = \mu(F_1) - \alpha.$$

□

The theorem is then completely proven. □

4.2.3 Determinant line bundles for triples

Let S be a connected scheme of finite type over k , $p : S \times X \rightarrow S$ and $q : S \times X \rightarrow X$ be the projections. Let $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ be a family of triples on X parametrized by S . For any triple $F = (F_2 \xrightarrow{\psi} F_1)$ on X , the triple product of \mathcal{T} and q^*F ,

$$\mathcal{T} \times q^*F = (\mathcal{E}_2 \otimes q^*F_1 \oplus \mathcal{E}_1 \otimes q^*F_2 \xrightarrow{\varphi_S \times q^*\psi} \mathcal{E}_1 \otimes q^*F_1),$$

is again a S -family of triples.

Definition 4.2.5. Let $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ be a family of triples on X parametrized by S and $F = (F_2 \xrightarrow{\psi} F_1)$ a triple on X . Then the line bundle

$$\mathcal{L}_{\mathcal{T} \times q^*F} := \lambda_{\mathcal{E}_1}([F_2]) \otimes \lambda_{\mathcal{E}_2}([F_1]) \otimes (\lambda_{\mathcal{E}_1}([F_1]))^{-1},$$

where $\lambda_{\mathcal{E}_i}$ are morphisms defined in (4.1), is called the determinant line bundle associated to \mathcal{T} at F . As \mathcal{T} is fixed, we denote $\mathcal{L}_{\mathcal{T} \times q^*F}$ just by $\mathcal{L}_{F_2 \rightarrow F_1}$.

We will show that the determinant line bundle associated to \mathcal{T} at F defined above has similar properties of the determinant line bundle associated to a family of vector bundles as in Lemma 4.1.1: let $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$, $\mathcal{T}' = (\mathcal{E}'_2 \xrightarrow{\varphi'_S} \mathcal{E}'_1)$ and $\mathcal{T}'' = (\mathcal{E}''_2 \xrightarrow{\varphi''_S} \mathcal{E}''_1)$ be S -families of triples on X such that the following sequence

$$0 \longrightarrow \mathcal{T}' \longrightarrow \mathcal{T} \longrightarrow \mathcal{T}'' \longrightarrow 0$$

is exact. Since the triple product is exact, the sequence

$$0 \longrightarrow \mathcal{T}' \times q^*F \longrightarrow \mathcal{T} \times q^*F \longrightarrow \mathcal{T}'' \times q^*F \longrightarrow 0$$

is exact. Precisely, we have the following commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{E}'_2 \otimes q^*F_1 \oplus \mathcal{E}'_1 \otimes q^*F_2 & \longrightarrow & \mathcal{E}_2 \otimes q^*F_1 \oplus \mathcal{E}_1 \otimes q^*F_2 & \longrightarrow & \mathcal{E}''_2 \otimes q^*F_1 \oplus \mathcal{E}''_1 \otimes q^*F_2 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{E}'_1 \otimes q^*F_1 & \longrightarrow & \mathcal{E}_1 \otimes q^*F_1 & \longrightarrow & \mathcal{E}''_1 \otimes q^*F_1 \longrightarrow 0, \end{array} \quad (4.18)$$

where the rows are exact. From the exactness of the rows in (4.18) we have

$$\lambda_{\mathcal{E}_1}([F_2]) \otimes \lambda_{\mathcal{E}_2}([F_1]) \cong \lambda_{\mathcal{E}'_1}([F_2]) \otimes \lambda_{\mathcal{E}'_2}([F_1]) \otimes \lambda_{\mathcal{E}''_1}([F_2]) \otimes \lambda_{\mathcal{E}''_2}([F_1]),$$

$$\lambda_{\mathcal{E}_1}([F_1]) \cong \lambda_{\mathcal{E}'_1}([F_1]) \otimes \lambda_{\mathcal{E}''_1}([F_1]).$$

It is implied by the definition that

$$\begin{aligned} \mathcal{L}_{\mathcal{T} \times q^*F} &:= \lambda_{\mathcal{E}_1}([F_2]) \otimes \lambda_{\mathcal{E}_2}([F_1]) \otimes (\lambda_{\mathcal{E}_1}([F_1]))^{-1} \\ &\cong \lambda_{\mathcal{E}'_1}([F_2]) \otimes \lambda_{\mathcal{E}'_2}([F_1]) \otimes \lambda_{\mathcal{E}''_1}([F_2]) \otimes \lambda_{\mathcal{E}''_2}([F_1]) \otimes \lambda_{\mathcal{E}'_1}([F_1])^{-1} \otimes \lambda_{\mathcal{E}''_1}([F_1])^{-1} \\ &\cong [\lambda_{\mathcal{E}'_1}([F_2]) \otimes \lambda_{\mathcal{E}'_2}([F_1]) \otimes \lambda_{\mathcal{E}'_1}([F_1])^{-1}] \otimes [\lambda_{\mathcal{E}''_1}([F_2]) \otimes \lambda_{\mathcal{E}''_2}([F_1]) \otimes \lambda_{\mathcal{E}''_1}([F_1])^{-1}] \\ &\cong \mathcal{L}_{\mathcal{T}' \times q^*F} \otimes \mathcal{L}_{\mathcal{T}'' \times q^*F}. \end{aligned}$$

Similarly, if $0 \longrightarrow F' \longrightarrow F \longrightarrow F'' \longrightarrow 0$ is an exact sequence of triples, we have

$$\mathcal{L}_{\mathcal{T} \times q^*F} \cong \mathcal{L}_{\mathcal{T} \times q^*F'} \otimes \mathcal{L}_{\mathcal{T} \times q^*F''}.$$

Using the distributive property of pullback over tensor product, the base-change property is also fulfilled, i.e. if $f : S' \longrightarrow S$ be a morphism of scheme then

$$f^* \mathcal{L}_{\mathcal{T} \times q^*F} = \mathcal{L}_{f_X^* \mathcal{T} \times q'^*F},$$

where $f_X = f \times \text{Id}_X : S' \times X \longrightarrow S \times X$ and $q' : S' \times X \longrightarrow X$ is the second projection. For any line bundle L on S , we have

$$\mathcal{T} \otimes p^*L = (\mathcal{E}_2 \otimes p^*L \xrightarrow{\varphi_S \times p^* \text{Id}_L} \mathcal{E}_1 \otimes p^*L)$$

and then

$$\begin{aligned}
\mathcal{L}_{\mathcal{T} \otimes p^* L \times q^* F} &= \lambda_{\mathcal{E}_1 \otimes p^* L}([F_2]) \otimes \lambda_{\mathcal{E}_2 \otimes p^* L}([F_1]) \otimes (\lambda_{\mathcal{E}_1 \otimes p^* L}([F_1]))^{-1} \\
&= \mathcal{L}_{\mathcal{T} \times q^* F} \otimes L^{\chi(\mathcal{E}_{1s} \otimes F_2) + \chi(\mathcal{E}_{2s} \otimes F_1) - \chi(\mathcal{E}_{1s} \otimes F_1)} \\
&= \mathcal{L}_{\mathcal{T} \times q^* F} \otimes L^{\chi(\mathcal{E}_s \times F)}
\end{aligned}$$

for some $s \in S$.

Since $\lambda_{\mathcal{E}_i}([F])$ depends only on the class $[F] \in K(X)$ (see (4.9)), we get the following similar property for determinant line bundles for triples.

Lemma 4.2.6. *Let $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ be a family of triples on X parametrized by S and $F = (F_2 \xrightarrow{\psi} F_1)$, $F' = (F'_2 \xrightarrow{\psi'} F'_1)$ be triples on X . If $\text{rank}(F'_i) = \text{rank}(F_i)$ and $\det(F'_i) = \det(F_i)$ for $i = 1, 2$ then*

$$\mathcal{L}_{\mathcal{T} \times q^* F'} \cong \mathcal{L}_{\mathcal{T} \times q^* F}.$$

4.2.4 Invariant sections

Let $F = (F_2 \xrightarrow{\psi} F_1)$ be a triple on X where ψ is surjective and $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ be a family of triples on X parametrized by S . By taking the triple product, we have an exact sequence of vector bundles on $S \times X$,

$$0 \longrightarrow \mathcal{K}_F \longrightarrow \mathcal{E}_2 \otimes q^* F_1 \oplus \mathcal{E}_1 \otimes q^* F_2 \longrightarrow \mathcal{E}_1 \otimes q^* F_1 \longrightarrow 0. \quad (4.19)$$

Hence

$$\mathcal{L}_{F_2 \rightarrow F_1} := \lambda_{\mathcal{E}_1}([F_2]) \otimes \lambda_{\mathcal{E}_2}([F_1]) \otimes (\lambda_{\mathcal{E}_1}([F_1]))^{-1} \cong \lambda_{\mathcal{K}_F}([\mathcal{O}_X]), \quad (4.20)$$

where $\lambda_{\mathcal{K}_F}([\mathcal{O}_X])$ is the determinant line bundle associated to S -family of vector bundles \mathcal{K}_F . Let us set $\theta_{F_2 \rightarrow F_1} := \mathcal{L}_{F_2 \rightarrow F_1}^{-1}$. Then $\theta_{F_2 \rightarrow F_1} \cong \lambda_{\mathcal{K}_F}([\mathcal{O}_X])^{-1}$. For any $s \in S$, we have an exact sequence of vector bundles on X

$$0 \longrightarrow (\mathcal{K}_F)_s \longrightarrow \mathcal{E}_{2s} \otimes F_1 \oplus \mathcal{E}_{1s} \otimes F_2 \longrightarrow \mathcal{E}_{1s} \otimes F_1 \longrightarrow 0.$$

It follows that $\chi((\mathcal{K}_F)_s) = \chi(\mathcal{E}_s \times F)$. As we have seen, if $\chi((\mathcal{K}_F)_s) = 0$ for all $s \in S$, the line bundle $\lambda_{\mathcal{K}_F}([\mathcal{O}_X])^{-1}$ has a geometric section, denoted by $\zeta_{\mathcal{K}_F}$. Hence $\theta_{F_2 \rightarrow F_1}$ also has a geometric section, denoted by $\zeta_{F_2 \rightarrow F_1}$, via the isomorphism $\theta_{F_2 \rightarrow F_1} \cong \lambda_{\mathcal{K}_F}([\mathcal{O}_X])^{-1}$. The zero divisor of $\zeta_{F_2 \rightarrow F_1}$, denoted by $\Theta_{F_2 \rightarrow F_1}$, parametrizes the same points as the zero divisor of $\zeta_{\mathcal{K}_F}$,

$$\Theta_{F_2 \rightarrow F_1} = \{s \in S \mid H^0(X, (\mathcal{K}_F)_s) \neq 0\}. \quad (4.21)$$

Let $T = (E_2 \xrightarrow{\varphi} E_1)$ be any triple of type (r_1, r_2, d_1, d_2) and $\alpha = \frac{a}{b}$ be a rational number. Let E_T be the extension bundle of rank $r = r_1 + r_2$ defined by T and $m_0 = \lceil \frac{1-r}{r} \Delta(E_T) + 1 \rceil$.

Let us set $d = m_0 \deg_{H(\alpha)}(E_T)$ and $h = \gcd(r, d)$. For each positive integer p , let $\text{Comp}(p)$ be the set of triples $F = (F_2 \xrightarrow{\psi} F_1)$ which have the following numerical invariants

$$\begin{aligned} \text{rank}(F_1) &= pm_0 br_0, \deg(F_1) = p(m_0 br_0(g_X - 1) - d_0), \\ \text{rank}(F_2) &= 2 \text{rank}(F_1), \deg(F_2) = 2 \deg(F_1) - \alpha \text{rank}(F_1). \end{aligned} \quad (4.22)$$

where $r_0 = \frac{r}{h}$, $d_0 = \frac{d}{h}$. These triples can be constructed as in the proof of Lemma 4.2.4. Let us set

$$\text{Comp} = \cup_{p \in \mathbb{Z}_{>0}} \text{Comp}(p). \quad (4.23)$$

In particular, for any $F \in \text{Comp}$ and any triple T of type (r_1, r_2, d_1, d_2) , we have

$$\begin{aligned} \chi(T \times F) &= \chi(E_2 \otimes F_1 \oplus E_1 \otimes F_2) - \chi(E_1 \otimes F_1) \\ &= (d_1 + d_2 - 2\alpha r_1) \text{rank}(F_1) + \deg(F_1)(r_1 + r_2) + r_{F_1}(r_1 + r_2)(1 - g_X) \\ &= (d_1 + d_2 - 2\alpha r_1)m_0 bpr_0 + r(m_0 bpr_0(g_X - 1) - pd_0) + rm_0 bpr_0(1 - g_X) \\ &= (d_1 + d_2 - 2\alpha r_1)m_0 bpr_0 - rpd_0 = 0 \end{aligned} \quad (4.24)$$

since $m_0 b(d_1 + d_2 - 2\alpha r_1) = m_0 \deg_{H(\alpha)}(E_T)$.

Assume that $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ is a S -family of triples of type (r_1, r_2, d_1, d_2) . By the definition of α -orthogonal triples, for any $F = (F_2 \xrightarrow{\psi} F_1) \in \text{Comp}$, the zero divisor of geometric section $\zeta_{F_2 \rightarrow F_1}$ is described as follows

$$\Theta_{F_2 \rightarrow F_1} = \{s \in S \mid F \text{ is not a } \alpha\text{-orthogonal triple of } \mathcal{T}_s\}. \quad (4.25)$$

4.3 Base point freeness

4.3.1 Generalized theta line bundles for triples

Let $T = (E_2 \xrightarrow{\varphi} E_1)$ be a α -semistable triple of type (r_1, r_2, d_1, d_2) on X where $\alpha = \frac{a}{b}$ and E_T be the extension on $X \times \mathbb{P}^1$ defined by T . We know that E_T is $H(\alpha)$ -semistable. Moreover, the restriction of E_T to smooth projective curve $C' \in |mH(\alpha)|$ is still semistable for any $m \geq m_0 = \lceil \frac{1-r}{r} \Delta(E_T) + 1 \rceil$. It is easy to see that T and T' are S -equivalent α -semistable triples if and only if E_T and $E_{T'}$, the extensions defined by T and T' , respectively, are S -equivalent. We will show that for $m \gg m_0$, the S -equivalence is preserved under the restriction to a smooth projective curve $C' \in |mH(\alpha)|$.

Lemma 4.3.1. *Let E_T and $E_{T'}$ be $H(\alpha)$ -semistable vector bundles on $X \times \mathbb{P}^1$ defined by triples T and T' , respectively. Then for m large enough, $E_T|_{C'}$ and $E_{T'}|_{C'}$ are S -equivalent if and only if T and T' are S -equivalent.*

Proof. Assume that

$$0 \subset T^1 \subset T^2 \subset \dots \subset T^n = T$$

is a Jordan-Hölder filtration of T . By the construction of extensions, it follows that

$$0 \subset E_{T^1} \subset E_{T^2} \subset \dots \subset E_{T^n} = E_T$$

is a Jordan-Hölder filtration of E_T . Since $E_{T^i}/E_{T^{i-1}}$ are stable of $H(\alpha)$ -slope $\mu_{H(\alpha)}(E_T)$ for all $i \geq 1$, for any $m \geq m_0$, $E_{T^i}|_{C'}/E_{T^{i-1}}|_{C'}$ are stable of the same slope (see also the proof of Corollary 2.3.4) since

$$E_{T^i}|_{C'}/E_{T^{i-1}}|_{C'} \cong (E_{T^i}/E_{T^{i-1}})|_{C'}.$$

Hence

$$0 \subset E_{T^1}|_{C'} \subset E_{T^2}|_{C'} \subset \dots \subset E_{T^n}|_{C'} = E_T|_{C'}$$

is again a Jordan-Hölder filtration of $E_T|_{C'}$. So we have

$$\text{gr}(E_T|_{C'}) \cong \text{gr}(E_T)|_{C'}.$$

It implies that $\text{gr}(E_T|_C) \cong \text{gr}(E_{T'}|_C)$ if $\text{gr}(T) \cong \text{gr}(T')$.

Conversely, it is enough to show that for m large enough, if E and E' are $H(\alpha)$ -stable vector bundles on $X \times \mathbb{P}^1$ of the same slope such that $E|_{C'}$ and $E'|_{C'}$ are isomorphic and stable then $E \cong E'$. Since E and E' are stable of the same slope, then

$$E \cong E' \iff H^0(X \times \mathbb{P}^1, E^\vee \otimes E') \neq 0.$$

Consider the following exact sequence on $X \times \mathbb{P}^1$,

$$0 \longrightarrow E^\vee \otimes E' \otimes \mathcal{O}_{X \times \mathbb{P}^1}(-mH(\alpha)) \longrightarrow E^\vee \otimes E' \longrightarrow (E^\vee \otimes E')|_{C'} \longrightarrow 0.$$

We then obtain a long exact sequence of cohomology

$$\begin{aligned} 0 &\longrightarrow H^0(X \times \mathbb{P}^1, E^\vee \otimes E' \otimes \mathcal{O}_{X \times \mathbb{P}^1}(-mH(\alpha))) \longrightarrow H^0(X \times \mathbb{P}^1, E^\vee \otimes E') \\ &\longrightarrow H^0(X \times \mathbb{P}^1, (E^\vee \otimes E')|_{C'}) \longrightarrow H^1(X \times \mathbb{P}^1, E^\vee \otimes E' \otimes \mathcal{O}_{X \times \mathbb{P}^1}(-mH(\alpha))) \end{aligned}$$

where $H^0(X \times \mathbb{P}^1, (E^\vee \otimes E')|_{C'}) \cong H^0(C', E^\vee|_{C'} \otimes E'|_{C'}) \neq 0$ since $E|_{C'} \cong E'|_{C'}$. We have $mH(\alpha) = maF_p + mbF_q$ and $a, b > 0$. As

$$E = E_T \in \text{Ext}^1(p^*E_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2), p^*E_2), E' = E_{T'} \in \text{Ext}^1(p^*E'_1 \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-2), p^*E'_2),$$

by Künneth formula, the vanishing of

$$H^0(X \times \mathbb{P}^1, E^\vee \otimes E' \otimes \mathcal{O}_{X \times \mathbb{P}^1}(-mH(\alpha))) \text{ and } H^1(X \times \mathbb{P}^1, E^\vee \otimes E' \otimes \mathcal{O}_{X \times \mathbb{P}^1}(-mH(\alpha)))$$

is induced from the vanishing of

$$H^0(X, E_j^\vee \otimes E'_i(-ma)) \text{ and } H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-mb+2))$$

which is satisfied for $m \gg m_0$. It follows that

$$H^0(X \times \mathbb{P}^1, E^\vee \otimes E') \cong H^0(X \times \mathbb{P}^1, (E^\vee \otimes E')|_{C'}) \neq 0.$$

Hence $E \cong E'$. □

This lemma implies that for any $m \geq m_0$, we have a well defined morphism between moduli spaces,

$$\begin{aligned} \mathbf{t}_m : M_\alpha^{ss}(\underline{r}, \underline{d}) &\longrightarrow U_{C'}(r, d') \\ T &\mapsto E_T|_{C'}. \end{aligned} \tag{4.26}$$

and \mathbf{t}_m is an injective if $m \gg m_0$.

Let F_C be a complementary vector bundle of $U_C(r, d)$ where $C \in |m_0 H(\alpha)|$. Then the generalized theta line bundle θ_{F_C} has a section ζ_{F_C} whose zero divisor is Θ_{F_C} . Let us set

$$\mathbf{t} = \mathbf{t}_{m_0} : M_\alpha^{ss}(\underline{r}, \underline{d}) \longrightarrow U_C(r, d).$$

Definition 4.3.2. *We call the pullback $\mathbf{t}^* \theta_{F_C}$, $\mathbf{t}^* \zeta_{F_C}$ again a generalized theta line bundle and a generalized theta function, respectively, on $M_\alpha^{ss}(\underline{r}, \underline{d})$.*

4.3.2 Linear systems

Assume that $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ is a family of α -semistable triples of type (r_1, r_2, d_1, d_2) on X parametrized by a connected scheme S . For any $F = (F_2 \xrightarrow{\psi} F_1) \in \text{Comp}$, the line bundle $\theta_{F_2 \rightarrow F_1}$ has a geometric section $\zeta_{F_2 \rightarrow F_1}$ whose zero divisor is

$$\Theta_{F_2 \rightarrow F_1} = \{s \in S \mid F = (F_2 \xrightarrow{\psi} F_1) \text{ is not a } \alpha - \text{orthogonal triple of } \mathcal{T}_s\}.$$

For simplicity, we denote by \mathcal{M} the moduli space $\mathcal{M}_\alpha^{ss}(\underline{r}, \underline{d})$. Consider the following diagram

$$S \xrightarrow{f_{\mathcal{T}}} \mathcal{M} \xrightarrow{\mathbf{t}} U_C(r, d)$$

where $f_{\mathcal{T}}$ is induced from family \mathcal{T} . Let us recall the linear systems which exist on these schemes:

+ On $U_C(r, d)$: we have pluritheta linear series $|p\Theta_{F_0}|$ where Θ_{F_0} is a basic theta divisor. For any complementary vector bundle F_C such that $\det(F_C) \cong L_p$, the generalized theta line bundle θ_{F_C} has a section ζ_{F_C} whose zero divisor is Θ_{F_C} . Moreover, θ_{F_C} is isomorphic to $\theta_{F_0}^p$ for such any F_C .

+ On \mathcal{M} : we set $\mathcal{L} = \mathfrak{t}^*\theta_{F_0}$ and call it a basic theta line bundle. For each p , we have a generalized theta line bundle \mathcal{L}^p and generalized theta functions $\mathfrak{t}^*(\zeta_{F_C})$ for any such complementary vector bundles F_C . Let $V(\mathcal{L}^p) \subset H^0(\mathcal{M}, \mathcal{L}^p)$ be the linear space spanned by sections $\mathfrak{t}^*\zeta_{F_C}$.

+ On S : Let us fix F_C a complementary vector bundle F_C of $U_C(r, d)$ as above, we consider the following triple constructed as in the proof of Lemma 4.2.4,

$$F = (F_2 \xrightarrow{\psi} F_1) := (f_*(V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)|_C \otimes F_C) \xrightarrow{\psi} f_*F_C), \quad (4.27)$$

where f is the composed morphism $C \hookrightarrow X \times \mathbb{P}^1 \rightarrow X$ and $q : X \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$ is the projection. It implies that $F \in \text{Comp}(p)$ and then $\chi(\mathcal{E}_s \times F) = 0$ for any $s \in S$. Let $\text{Comp}(p, F) \subset \text{Comp}(p)$ be the subset of triples $F' = (F'_2 \xrightarrow{\psi'} F'_1) \in \text{Comp}(p)$ such that $[F'_i] = [F_i] \in K(X), i = 1, 2$. It follows from Lemma 4.2.6 that all line bundles $\theta_{F'_2 \rightarrow F'_1}$ are isomorphic to $\theta_{F_2 \rightarrow F_1}$. Hence we obtain a unique line bundle $\theta_{F_2 \rightarrow F_1}$ and a linear system $V(p, F) \subset H^0(S, \theta_{F_2 \rightarrow F_1})$ spanned by geometric sections $\zeta_{F'_2 \rightarrow F'_1}$ where $F' = (F'_2 \xrightarrow{\psi'} F'_1) \in \text{Comp}(p, F)$.

4.3.3 Corollaries

The separation property of $|p\Theta_{F_0}|$ on $U_C(r, d)$ implies that for any semistable vector bundle $E_C \in U_C(r, d)$ and any $p \geq p_0 := r^2 + r$, there exists a complementary vector bundle F_C of determinant L_p such that $\Theta_{F_C} \in |p\Theta_{F_0}|$ and $E_C \notin \text{Supp } \Theta_{F_C}$. As a consequence, we obtain similar result for the base point freeness of the linear systems mentioned above on S and \mathcal{M} .

For any triple $F' = (F'_2 \xrightarrow{\psi'} F'_1) \in \text{Comp}(p, F)$, we set

$$\overline{\Theta}_{F'_2 \rightarrow F'_1} := \{T \in \mathcal{M} | F' = (F'_2 \xrightarrow{\psi'} F'_1) \text{ is not a } \alpha\text{-orthogonal triple of } T\}.$$

Corollary 4.3.3. *For any $p \geq p_0$, then*

(i) *The linear systems $V(\mathcal{L}^p)$ and $V(p, F)$ are base point free.*

(ii) *There exists a triple $F' = (F'_2 \xrightarrow{\psi'} F'_1) \in \text{Comp}(p, F)$ such that $\overline{\Theta}_{F'_2 \rightarrow F'_1}$ is a Cartier divisor on \mathcal{M} and*

$$\mathcal{L}^p \cong \mathcal{O}_{\mathcal{M}}(\overline{\Theta}_{F'_2 \rightarrow F'_1}).$$

Proof. Let T be any α -semistable triple on X . For the proof, we may assume that there exists $s \in S$ such that $\mathcal{E}_s = T$. Let E_T be the extension defined by T on $X \times \mathbb{P}^1$. Then $\mathfrak{t}(T) = E_T|_C \in U_C(r, d)$. Since $|p\Theta_{F_0}|$ is base point free, there exists a complementary vector

bundle F'_C such that $\det(F'_C) \cong L_p$, $\Theta_{F'_C} \in |p\Theta_{F_0}|$ and $E_T|_C \notin \text{Supp } \Theta_{F'_C}$. It implies that $\mathfrak{t}^*(\zeta_{F'_C})(T) \neq 0$. Let

$$F' = (F'_2 \xrightarrow{\psi'} F'_1) := (f_*(V \otimes q^*\mathcal{O}_{\mathbb{P}^1}(-1)|_C \otimes F'_C) \xrightarrow{\psi'} f_*F'_C),$$

be the triple constructed from F'_C . Then $F' \in \text{Comp}(p, F)$ and moreover, it is a α -orthogonal triple of $T = \mathcal{E}_s$. Therefore $\zeta_{F'_2 \rightarrow F'_1}(s) \neq 0$. This proves (i) and moreover implies that $\mathfrak{t}^*\Theta_{F'_C}$ is a Cartier divisor. In particular, we have

$$\begin{aligned} \mathfrak{t}^*\Theta_{F'_C} &= \{T \in \mathcal{M} | E_T|_C \in \Theta_{F'_C}\} \\ &= \{T \in \mathcal{M} | H^0(C, E_T|_C \otimes F'_C) \neq 0\} \\ &= \{T \in \mathcal{M} | F' = (F'_2 \xrightarrow{\psi'} F'_1) \text{ is not a } \alpha\text{-orthogonal triple of } T\} \\ &= \overline{\Theta}_{F'_2 \rightarrow F'_1}. \end{aligned} \tag{4.28}$$

It follows that

$$\mathcal{L}^p = \mathfrak{t}^*\theta_{F_0}^p \cong \mathfrak{t}^*\mathcal{O}_{U_C(r,d)}(\Theta_{F'_C}) \cong \mathcal{O}_{\mathcal{M}}(\overline{\Theta}_{F'_2 \rightarrow F'_1}).$$

□

By the construction of the extensions defined by triples, for each S -family of α -semistable triples $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ we obtain a family $E_{\mathcal{T}} = (E_{\mathcal{T}_s})_{s \in S}$ of $H(\alpha)$ -semistable extensions on $X \times \mathbb{P}^1$ parametrized by S . Restricting this family to C , we get a family of semistable vector bundles on C ,

$$\mathcal{E} = (E_{\mathcal{T}_s}|_C)_{s \in S}.$$

We also have a commutative diagram

$$\begin{array}{ccc} S & \xrightarrow{f_{\mathcal{E}}} & U_C(r, d) \\ & \searrow f_{\mathcal{T}} & \nearrow \mathfrak{t} \\ & \mathcal{M} & \end{array} \tag{4.29}$$

where $f_{\mathcal{E}}$ is the morphism induced by \mathcal{E} . For any complementary vector bundle F_C of $U_C(r, d)$, it is implied by Theorem 4.1.2 that the line bundle $\lambda_{\mathcal{E}}([F_C])^{-1}$ descends to θ_{F_C} by $f_{\mathcal{E}}$. We obtain a similar universal property for the generalized theta line bundles on \mathcal{M} .

Corollary 4.3.4. *Let $p \geq 1$ and $F'' = (F''_2 \xrightarrow{\psi''} F''_1)$ be any triple in $\text{Comp}(p, F)$. Then for any family $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_S} \mathcal{E}_1)$ of α -semistable triples on X parametrized by S , the line bundle $\theta_{F''_2 \rightarrow F''_1}$ descends to \mathcal{L}^p .*

Proof. Since $f_{\mathcal{E}} = \mathbf{t} \circ f_{\mathcal{T}}$, $\lambda_{\mathcal{E}}([F_C])^{-1} \cong f_{\mathcal{T}}^* \mathcal{L}^p$ for any complementary vector bundle F_C such that $\det(F_C) = L_p$. Consider the following diagram

$$\begin{array}{ccccc}
 S \times C & \xrightarrow{q_C} & C & & \\
 \swarrow p_C & \searrow i_S & \downarrow & \searrow i & \\
 S & & S \times X \times \mathbb{P}^1 & \xrightarrow{f} & X \times \mathbb{P}^1 \xrightarrow{q} \mathbb{P}^1 \\
 \nwarrow p_X & \downarrow f_S & \nwarrow p_{12} & \downarrow p_{23} & \nwarrow p \\
 & S \times X & \xrightarrow{q_X} & X &
 \end{array} \tag{4.30}$$

By the construction of $E_{\mathcal{T}}$, we have an exact sequence

$$0 \longrightarrow E_{\mathcal{T}} \longrightarrow p_{12}^* \mathcal{E}_2 \oplus p_{12}^* \mathcal{E}_1 \otimes V \otimes (q \circ p_{23})^* \mathcal{O}_{\mathbb{P}^1}(-1) \longrightarrow p_{12}^* \mathcal{E}_1 \longrightarrow 0.$$

Restricting to $S \times C$, we have

$$0 \longrightarrow \mathcal{E} \longrightarrow p_{12}^* \mathcal{E}_2|_{S \times C} \oplus p_{12}^* \mathcal{E}_1 \otimes V \otimes (q \circ p_{23})^* \mathcal{O}_{\mathbb{P}^1}(-1)|_{S \times C} \longrightarrow p_{12}^* \mathcal{E}_1|_{S \times C} \longrightarrow 0$$

(see (3.12) and (3.11)). By twisting with $q_C^* F_C$, we pass to the following exact sequence

$$\begin{aligned}
 0 \longrightarrow \mathcal{E} \otimes q_C^* F_C &\longrightarrow p_{12}^* \mathcal{E}_2|_{S \times C} \otimes q_C^* F_C \oplus p_{12}^* \mathcal{E}_1 \otimes V \otimes (q \circ p_{23})^* \mathcal{O}_{\mathbb{P}^1}(-1)|_{S \times C} \otimes q_C^* F_C \\
 &\longrightarrow p_{12}^* \mathcal{E}_1|_{S \times C} \otimes q_C^* F_C \longrightarrow 0.
 \end{aligned} \tag{4.31}$$

Since f is finite morphism, $(f_S)_*$ is an exact functor. We then obtain a short exact sequence on $S \times X$,

$$\begin{aligned}
 0 &\longrightarrow (f_S)_*(\mathcal{E} \otimes q_C^* F_C) \\
 &\longrightarrow (f_S)_*(p_{12}^* \mathcal{E}_2|_{S \times C} \otimes q_C^* F_C) \oplus (f_S)_*(p_{12}^* \mathcal{E}_1 \otimes V \otimes (q \circ p_{23})^* \mathcal{O}_{\mathbb{P}^1}(-1)|_{S \times C} \otimes q_C^* F_C) \\
 &\longrightarrow (f_S)_*(p_{12}^* \mathcal{E}_1|_{S \times C} \otimes q_C^* F_C) \longrightarrow 0.
 \end{aligned} \tag{4.32}$$

By projection formula and flat base change, we obtain the following exact sequence

$$0 \longrightarrow (f_S)_*(\mathcal{E} \otimes q_C^* F_C) \longrightarrow \mathcal{E}_2 \otimes q_X^* F_1 \oplus \mathcal{E}_1 \otimes q_X^* F_2 \longrightarrow \mathcal{E}_1 \otimes q_X^* F_1 \longrightarrow 0.$$

Since $p_C = p_X \circ f_S$ and $(f_S)_*$ is exact,

$$\theta_{F_2 \rightarrow F_1} \cong \lambda_{(f_S)_*(\mathcal{E} \otimes q_C^* F_C)}([\mathcal{O}_X])^{-1} \cong \lambda_{\mathcal{E}}([F_C])^{-1}.$$

Now for any $F'' = (F_2'' \xrightarrow{\psi''} F_1'') \in \text{Comp}(p, F)$,

$$\theta_{F_2'' \rightarrow F_1''} \cong \theta_{F_2 \rightarrow F_1} \cong f_{\mathcal{T}}^* \mathcal{L}^p.$$

□

We consider the numerical equivalence on $K(X \times \mathbb{P}^1)$ which generalizes the numerical equivalence of divisors defined in Chapter 2 (cf. [12, §8, p.178]). Let u and u' be classes in $K(X \times \mathbb{P}^1)$. Then they are said to be numerically equivalent: $u \equiv u'$, if their difference is contained in the radical of the quadratic form $(a, b) \mapsto \chi(a \cdot b)$. Let

$$K(X \times \mathbb{P}^1)_{\text{Num}} = K(X \times \mathbb{P}^1) / \equiv.$$

It is easy to see that for any triple T of fixed type (r_1, r_2, d_1, d_2) , the corresponding extension E_T is contained in a fixed class $\mathfrak{e} \in K(X \times \mathbb{P}^1)_{\text{Num}}$. Let $U_{X \times \mathbb{P}^1}(\mathfrak{e})$ be the moduli space of $H(\alpha)$ -semistable vector bundles on $X \times \mathbb{P}^1$ of numerical class \mathfrak{e} .

Corollary 4.3.5. *The line bundle \mathcal{L} on \mathcal{M} is ample.*

Proof. It is enough to show that, \mathcal{L}^p is ample for some $p \geq p_0$. Let F_C be a complementary vector bundle of $U_C(r, d)$ such that $\mathcal{L}^p = \mathfrak{t}^* \theta_{F_C}$. Let $C' \in |mC|$ where m large enough such that the morphism

$$\mathfrak{t}_m : \mathcal{M} \longrightarrow U_{C'}(r, md)$$

is an injection (see (4.26)). If we can find a complementary vector bundle $F_{C'}$ of $U_{C'}(r, md)$ such that $\mathfrak{t}_m^* \theta_{F_{C'}} \cong \mathcal{L}^{mp}$, we are done since $\mathfrak{t}_m^* \theta_{F_{C'}}$ is ample.

Assume that F_C has rank R and degree D . Consider the class $[F_C] \in K(X \times \mathbb{P}^1)$, we have

$$\text{ch}([F_C]) = R \text{ch}([\mathcal{O}_C]) + D = R.C + (D - \frac{RC^2}{2}).$$

Let K be the canonical divisor on $X \times \mathbb{P}^1$, then

$$g_C = \frac{1}{2}C^2 + \frac{1}{2}C.K + 1 \text{ and } g_{C'} = \frac{1}{2}C'^2 + \frac{1}{2}C'.K + 1.$$

As $C' \in |mC|$, we have

$$g_{C'} - 1 = m(g_C - 1) + \frac{m(m-1)}{2}C^2.$$

Let $F_{C'}$ be a vector bundle on C' of rank R and degree $D' = mD + \frac{m(m-1)}{2}RC^2$. We then have

$$rD' + Rmd + rR(1 - g_{C'}) = m(rD + Rd + rR(1 - g_C)) = 0,$$

i.e. $F_{C'}$ is a complementary vector bundle of $U_{C'}(r, md)$. Moreover,

$$\text{ch}([\mathcal{O}_{C'}]) = m \text{ch}([\mathcal{O}_C]) - \frac{m(m-1)}{2}C^2$$

and then

$$\text{ch}([F_{C'}]) = R \text{ch}([\mathcal{O}_{C'}]) + D' = R(m \text{ch}([\mathcal{O}_C]) - \frac{m(m-1)}{2}C^2) + D' = m \text{ch}([F_C]).$$

It implies that $[F_{C'}] = m[F_C] \in K(X \times \mathbb{P}^1)$.

Consider the following factors of \mathfrak{t} and \mathfrak{t}_m

$$\begin{array}{ccc}
 & U_C(r, d) & \\
 \mathfrak{t} \nearrow & \uparrow i^* & \\
 \mathcal{M} & \longrightarrow U_{X \times \mathbb{P}^1}(\mathfrak{e}) & \\
 \searrow \mathfrak{t}_m & \downarrow i_m^* & \\
 & U_{C'}(r, md) &
 \end{array} \tag{4.33}$$

where i^* and i_m^* are restrictions induced from the inclusions

$$i : C \hookrightarrow X \times \mathbb{P}^1 \text{ and } i_m : C' \hookrightarrow X \times \mathbb{P}^1,$$

respectively. Since $[F_{C'}] = m[F_C]$, it implies that $i_m^* \theta_{F_{C'}} \cong (i^* \theta_{F_C})^m$. Therefore, we have

$$\mathfrak{t}_m^* \theta_{F_{C'}} \cong (\mathfrak{t}^* \theta_{F_C})^m \cong \mathcal{L}^{mp}.$$

□

4.4 Langton's valuative criterion for triples

4.4.1 Elementary transformations

Let R be a discrete valuation ring where $\text{Spec } R = \{\eta, 0\}$ and $X_R = \text{Spec } R \times X$. Let us denote by $X_\eta = \eta \times X$ the generic fiber and X_0 the closed fiber of the first projection

$$p_R : \text{Spec } R \times X \longrightarrow \text{Spec } R.$$

Then X_0 is a Cartier divisor on X_R and any vector bundle on X_0 is a sheaf on X_R of projective dimension 1. Let \mathcal{E} be a vector bundle on X_R and E_0 be the restriction of \mathcal{E} on closed fiber X_0 . We recall now the construction of elementary transformations of \mathcal{E} along a quotient bundle of E_0 , from which we can similarly construct the elementary transformations of triples.

Let E be a vector bundle on X_0 which is a quotient of E_0 . Then the kernel \mathcal{E}' of the composition

$$\pi : \mathcal{E} \rightarrow E_0 \xrightarrow{\pi_0} E$$

is a vector bundle on X_R . We call it an elementary transformation of \mathcal{E} along E . If \mathcal{E}' is an elementary transformation of \mathcal{E} along E , we have a short exact sequence of sheaves on X_R

$$0 \rightarrow \mathcal{E}' \rightarrow \mathcal{E} \xrightarrow{\pi} E \rightarrow 0. \tag{4.34}$$

By restricting this exact sequence to X_η , we see easily that \mathcal{E}' and \mathcal{E} coincide over the generic fiber. On closed fiber, E becomes a subbundle of \mathcal{E}' .

Similarly, let $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_R} \mathcal{E}_1)$ be a triple on X_R and $T = (E_2 \xrightarrow{\varphi} E_1)$ be a quotient of T_0 , the restriction of \mathcal{T} on the closed fiber X_0 . Let $\phi : \mathcal{T} \rightarrow T$ be the composed morphism of triples on X_R .

Definition 4.4.1. *The triple $\mathcal{T}' = (\mathcal{E}'_2 \xrightarrow{\varphi'_R} \mathcal{E}'_1)$ is called an elementary transformation of \mathcal{T} along T if it satisfies the following short exact sequence of triples*

$$0 \rightarrow \mathcal{T}' \rightarrow \mathcal{T} \xrightarrow{\phi} T \rightarrow 0. \quad (4.35)$$

Notice that, the sequence (4.35) is a short exact sequence of triples of sheaves on X_R . But \mathcal{T}' is really a triple on X_R , i.e. \mathcal{E}'_1 and \mathcal{E}'_2 are vector bundle on X_R . As above, we can also conclude that the elementary transformations of \mathcal{T} along T do not change the triple over the generic point. But they transfer the quotient T of T_0 to a subtriple of \mathcal{T}' over the closed point.

Lemma 4.4.2. *Let $T = (E_2 \xrightarrow{\varphi} E_1)$ be a triple on X which is not α -semistable and $T' = (E'_2 \xrightarrow{\varphi'} E'_1)$ be the maximal destabilizing subtriple of T . Then the quotient*

$$T'' = T/T' := (E''_2 \xrightarrow{\varphi''} E''_1)$$

is again a triple, i.e. E''_i is vector bundle on X , $i = 1, 2$.

Proof. Let F'_i be the torsion subsheaf of E''_i , $i = 1, 2$. It is clear that $\varphi''(F'_2) \subset F'_1$. Then $F_i = E''_i/F'_i$ are vector bundles and $F = (F_2 \xrightarrow{\varphi''} F_1)$ is a triple. We have a short exact sequence

$$0 \longrightarrow F' \longrightarrow T'' \longrightarrow F \longrightarrow 0$$

where $F' = (F'_2 \xrightarrow{\varphi''} F'_1)$. Let G be the kernel of the composed morphism $T' \rightarrow T'' \rightarrow F$. We obtain a short exact sequence

$$0 \longrightarrow T' \longrightarrow G \longrightarrow F' \longrightarrow 0.$$

It implies that,

$$\mu_\alpha(G) = \mu_\alpha(T') + \frac{\text{length}(F'_1) + \text{length}(F'_2)}{r'_1 + r'_2} \geq \mu_\alpha(T'),$$

where r'_i is the rank of E'_i , $i = 1, 2$. Since T' is maximal destabilizing subtriple of T , $T' = G$. It follows that $F'_i = 0$ for $i = 1, 2$. Hence E''_i are vector bundles. \square

4.4.2 Valuative criterion

Lemma 4.4.3 ([10, Lm. 6.1]). *Let X be a smooth projective curve. Any vector bundle E on X_η can be extended to a vector bundle \mathcal{E} on X_R .*

In [10, Thm. 6.4], Hein gave a proof of Langton's theorem for semistable vector bundles of rank 2 and determinant ω_X using the elementary transformations and the cohomological characterization of semistability of vector bundles. By the same method, we can prove the following Langton's theorem for α -semistable triples (for the original Langton's valuative criterion, see [14]).

Theorem 4.4.4. *Let T_η be a α -semistable triple of type (r_1, r_2, d_1, d_2) on the generic fiber X_η . Then there exists a triple \mathcal{T} on X_R such that $\mathcal{T}|_{X_\eta} \cong T_\eta$ and $T_0 := \mathcal{T}|_{X_0}$ is α -semistable.*

Proof. Assume that $T_\eta = (E_{\eta 2} \xrightarrow{\varphi_\eta} E_{\eta 1})$ on X_η . It is implied from Lemma 4.4.3 that there exist vector bundles \mathcal{E}_2 and \mathcal{E}_1 on X_R which extend $E_{\eta 2}$ and $E_{\eta 1}$, respectively. We consider the morphism $\varphi_\eta \in \text{Hom}_{X_\eta}(E_{\eta 2}, E_{\eta 1})$ as a section of $\mathcal{E}_2^\vee \otimes \mathcal{E}_1$ over the open set X_η . By a standard argument as in [9, Lm. 5.14, p. 118], we can extend this section to obtain a section $\varphi_R \in \text{Hom}_{X_R}(\mathcal{E}_2, \mathcal{E}_1)$. Hence we get an extension $\mathcal{T} = (\mathcal{E}_2 \xrightarrow{\varphi_R} \mathcal{E}_1)$ of T_η on X_R .

Let $p_R : X_R \rightarrow \text{Spec } R$ and $q_R : X_R \rightarrow X$ be the projections. For any $F = (F_2 \xrightarrow{\psi} F_1) \in \text{Comp}$, we take the triple product $\mathcal{T} \times q_R^* F$ and obtain a short exact sequence of vector bundles on X_R ,

$$0 \longrightarrow \mathcal{K}_F \longrightarrow \mathcal{E}_2 \otimes q_R^* F_1 \oplus \mathcal{E}_1 \otimes q_R^* F_2 \longrightarrow \mathcal{E}_1 \otimes q_R^* F_1 \longrightarrow 0.$$

We now define an integer $\text{bad}(\mathcal{T})$

$$\text{bad}(\mathcal{T}) := \min_{F \in \text{Comp}} \{\text{bad}(\mathcal{K}_F)\}$$

where $\text{bad}(\mathcal{K}_F)$ is the number introduced first by Hein in [10]: it is equal to ∞ if $\text{Supp}(R^1 p_{R*} \mathcal{K}_F)$ is whole $\text{Spec } R$ and to $\text{length}(R^1 p_{R*} \mathcal{K}_F)$ otherwise.

Since T_η is α -semistable, it follows from Lemma 4.2.4 that there exists $F_\eta \in \text{Comp}$ such that $H^1(X_\eta, K_\eta) = 0$ where K_η is the restriction of \mathcal{K}_{F_η} to X_η . Therefore $R^1 p_{R*} \mathcal{K}_{F_\eta}$ is zero at generic point η . So we have

$$\text{bad}(\mathcal{T}) \leq \text{length}(R^1 p_{R*} \mathcal{K}_{F_\eta}),$$

which is finite. It is also implied by Theorem 4.2.3 and base change that

$$T_0 \text{ is } \alpha\text{-semistable} \iff \text{bad}(\mathcal{T}) = 0.$$

Assume that T_0 is not α -semistable. Then $\text{bad}(\mathcal{T})$ is positive. It is implied by Lemma 4.4.2 that there exists a quotient T of T_0 such that

$$\mu_\alpha(T) < \mu_\alpha(T_0).$$

Let \mathcal{T}' be an elementary transformation of \mathcal{T} along T . If we can show that

$$\text{bad}(\mathcal{T}') < \text{bad}(\mathcal{T})$$

then after taking a finite number of elementary transformations, we obtain the right one, i.e. an elementary transformation \mathcal{T}' such that $\text{bad}(\mathcal{T}')$ is zero.

For any $F \in \text{Comp}$, we have (see (4.24))

$$\chi(T_s \times F) = \chi(K_s) = 0$$

where T_s and K_s are the restriction of \mathcal{T} and \mathcal{K}_F , respectively, on the fiber over $s \in \text{Spec } R$. Choose a triple $F = (F_2 \xrightarrow{\psi} F_1) \in \text{Comp}$ such that

$$\text{bad}(\mathcal{T}) = \text{length}(R^1 p_{R*} \mathcal{K}_F).$$

It implies that $R^1 p_{R*} \mathcal{K}_F$ is supported only on the closed point $0 \in \text{Spec } R$. So we have

$$H^0(X_\eta, K_\eta) = H^1(X_\eta, K_\eta) = 0.$$

Hence $p_{R*} \mathcal{K}_F = 0$ since it is torsion free and zero at generic point.

Consider the following exact sequence of triples on X_R ,

$$0 \longrightarrow \mathcal{T}' \longrightarrow \mathcal{T} \longrightarrow T \longrightarrow 0.$$

By taking triple product with $q_R^* F$ and applying the kernel-cokernel sequence to the obtained diagram, we have the following exact sequence

$$0 \longrightarrow \mathcal{K}'_F \longrightarrow \mathcal{K}_F \longrightarrow K \longrightarrow 0.$$

Since $p_{R*} \mathcal{K}_F = 0$, by applying functor p_{R*} to this exact sequence, we get an exact sequence

$$0 \longrightarrow p_{R*} K \longrightarrow R^1 p_{R*} \mathcal{K}'_F \longrightarrow R^1 p_{R*} \mathcal{K}_F \longrightarrow R^1 p_{R*} K \longrightarrow 0.$$

We have $p_{R*} K = H^0(X_0, K)$ and $R^1 p_{R*} K = H^1(X_0, K)$ which are modules of finite length on R . Hence $R^1 p_{R*} \mathcal{K}'_F$ has finite length on R . In particular, we have

$$\text{length}(R^1 p_{R*} \mathcal{K}'_F) = \text{length}(R^1 p_{R*} \mathcal{K}_F) + \chi(K).$$

Assume that $T = (E_2 \xrightarrow{\varphi} E_1)$ on $X_0 \cong X$, then

$$0 \longrightarrow K \longrightarrow E_1 \otimes F_2 \oplus E_2 \otimes F_1 \longrightarrow E_1 \otimes F_1 \longrightarrow 0,$$

is an exact sequence. We have

$$\mu(K) = \mu_\alpha(T) + \mu(F_1)$$

(see also (4.14)). Similarly $\mu(K_0) = \mu_\alpha(T_0) + \mu(F_1)$. It implies that $\mu(K) < \mu(K_0)$ since $\mu_\alpha(T_0) > \mu_\alpha(T)$. So we have

$$\chi(K) < \chi(K_0) = 0$$

and then

$$\text{length}(R^1 p_{R*} \mathcal{K}'_F) < \text{length}(R^1 p_{R*} \mathcal{K}_F).$$

□

Bibliography

- [1] L. Álvarez-Cónsul. Some results on the moduli spaces of quiver bundles. *Geometriae Dedicata*, 139(1):99–120, 2009.
- [2] S. B. Bradlow and O. García-Prada. Stable triples, equivariant bundles and dimensional reduction. *Mathematische Annalen*, 304(2):225–252, 1996.
- [3] S. B. Bradlow, O. García-Prada, and P. B. Gothen. Moduli spaces of holomorphic triples over compact riemann surfaces. *Mathematische Annalen*, 328(1-2):299–351, 2004.
- [4] S. B. Bradlow, O. García-Prada, and P. B. Gothen. Moduli spaces of holomorphic triples over compact riemann surfaces. *Mathematische Annalen*, 328(1-2):299–351, 2004.
- [5] J.-M. Drezet and M. S. Narasimhan. Groupe de picard des variétés de modules de fibrés semi-stables sur les courbes algébriques. *Inventiones mathematicae*, 97(1):53–94, 1989.
- [6] E. Esteves. Separation properties of theta functions. *Duke Mathematical Journal*, 98(3):565–593, 1999.
- [7] E. Esteves and M. Popa. Effective very ampleness for generalized theta divisors. *Duke Mathematical Journal*, 123(3):429–444, 2004.
- [8] G. Faltings. Stable g-bundles and projective connections. *J. Algebraic Geom*, 2(3):507–568, 1993.
- [9] R. Hartshorne. Algebraic geometry, volume 52 of graduate texts in mathematics, 1977.
- [10] G. Hein. Duality construction of moduli spaces. *Geom. Dedicata*, 75(1):101–114, 1999.
- [11] G. Hein. Faltings’ construction of the moduli space of vector bundles on a smooth projective curve. In A. Schmitt, editor, *Affine Flag Manifolds and Principal Bundles*, Trends in Mathematics, pages 91–122. Springer Basel, 2010.
- [12] D. Huybrechts and M. Lehn. *The geometry of moduli spaces of sheaves*. Springer, 2010.

- [13] A. Langer. Semistable sheaves in positive characteristic. *Annals of mathematics*, pages 251–276, 2004.
- [14] S. G. Langton. Valuative criteria for families of vector bundles on algebraic varieties. *Annals of Mathematics*, pages 88–110, 1975.
- [15] J. Le Potier. Module des fibrés semi-stables et fonctions thêta. *Lecture Notes in Pure and Applied Mathematics*, pages 83–102, 1996.
- [16] M. Popa. Verlinde bundles and generalized theta linear series. *Transactions of the American Mathematical Society*, 354(5):1869–1898, 2002.
- [17] A. Schmitt. Moduli for decorated tuples of sheaves and representation spaces for quivers. In *Proceedings of the Indian Academy of Sciences-Mathematical Sciences*, volume 115, pages 15–49, 2005.
- [18] C. S. Seshadri. Vector bundles on curves. *Contemporary Mathematics*, 153:163, 1993.